INDEX TO AUTHORS

A

ABNEY, E. R. See KEARNEY, COOPER, KLEIN, ABNEY, PARKHOUSE, and LAWTON 297

ACTON, RONALD T. See ZWERNER, BARSTAD, and ACTON 986

AGUET, M. See ANDERSSON, AGUET, WIGHT, ANDERSSON, BINZ, and WIGZELL 1124

AHMED, AFTAB. See DICKLER, AHMED, and SACHS 1678

ALLISON, A. C. See HALSTEAD, O'ROURKE, and ALLISON 218

ANDERLE, SONIA K. See CROMARTIE, CRADDOCK, SCHWAB, ANDERLE, and YANG 1585

ANDERSSON, L. C., AGUET, M., WIGHT, E., ANDERSSON, R., BINZ, H., and WIGZELL, H. Induction of specific immune unresponsiveness using purified mixed leukocyte culture-activated T lymphoblasts as autoimmunogen. I. Demonstration of general validity as to species and histocompatibility barriers 1124

ANDERSSON, R. See ANDERSSON, AGUET, WIGHT, ANDERSSON, BINZ, and WIGZELL 1124

ASKENASE, PHILIP W. See CONE, GERSHON, and ASKENASE 1390

AUSTEN, K. FRANK. See FEARON and AUSTEN 22

—. See YURT and AUSTEN 1405

B

BANG, F. B. See WEISER and BANG 1467

BANKHURST, ARTHUR D. See GOODWIN, BANKHURST, and MESSNER 1719

BARSTAD, PAUL A. See ZWERNER, BARSTAD, and ACTON 986

BEAN, M. A., KODERA, Y., CUMMINGS, K. B., and BLOOM, B. R. Occurrence of restricted suppressor T-cell activity in man 1455

BECKER, CARL G., and DUBIN, THEODORE. Activation of factor XII by tobacco glycoprotein 457

BELLONE, C. J. See PRANGE, FIEDLER, NITECKI, and BELLONE 766

BENACERRAF, BARUJ. See BILLINGS, BURAKOFF, DORF, and BENACERRAF 623

—. See GREENE, PIERRES, DORF, and BENACERRAF 293

—. See PIERCE, GERMAIN, KAPP, and BENACERRAF 1827

—. See THÈZE, WALTENBAUGH, DORF, and BENACERRAF 287

—. See WALTENBAUGH, THÈZE, KAPP, and BENACERRAF 970

BENNINK, JACK. See BIDDISON, SNODGRASS, BENNINK, EFFROS, and DOHERTY 617

BERMAN, MONIQUE A., and WEIGLE, WILLIAM O. B-lymphocyte activation by the Fc region of IgG 241

BICK, PETER H., and MÖLLER, GÖRAN. Cytotoxic T-cell activation by polyribonucleotides: DNA synthesis is not required 844

—, PERSSON, ULLA, SMITH, EDVARD, MÖLLER, ERNA, and HAMMARSTRÖM, LENNART. Genetic control of lymphocyte activation: lack of response to low doses of concanavalin A in lipopolysaccharide-nonresponder mice 1146

BIDDISON, WILLIAM E., DOHERTY, PETER C., and WEBSTER, ROBERT G. Antibody to influenza virus matrix protein detects a common antigen on the surface of cells infected with type A influenza viruses 690

—, SNODGRASS, H. RALPH, BENNICK, JACK, EFFROS, RITA B., and DOHERTY, PETER C. Induction of virus-specific modifications recognized by cytotoxic T cells is not altered by prior substitution of target cells with trinitrophenyl 617

BILLING, RONALD. See CLINE and BILLING 1143

BILLINGS, PAUL, BURAKOFF, STEVEN, DORF, MARTIN E., and BENACERRAF, BARUJ. Cytotoxic T lymphocytes induced against allogeneic *I*-region determinants react with Ia molecules on trinitrophenyl-conjugated syngeneic target cells 623

BINZ, H. See ANDERSSON, AGUET, WIGHT, ANDERSSON, BINZ, and WIGZELL 1124

BLANDEN, ROBERT V., MCKENZIE, IAN F. C., KEES, URSULA, MELVOLD, ROGER W., and KOHN, HENRY I. Cytotoxic T-cell response to ectromelia virus-infected cells. Different *H-2* requirements for triggering precursor T-cell induction or lysis by effector T cells defined by the BALB/c-*H-2*^{db} mutation 869

BLOOM, BARRY R. See BEAN, KODERA, CUMMINGS, and BLOOM 1455

—. See SENIK and BLOOM 11

BONA, C., and CAZENAVE, P. A. Release from maternally-induced allotypic suppression in rabbit by Nocardia water-soluble mitogen 881

BOSMA, M. J., DE WITT, C., HAUSMAN, S. J., MARKS, R., and POTTER, M. Serological distinction of heavy chain variable regions (V_H subgroups) of mouse immunoglobulins. I. Common V_H determinants on the heavy chains of mouse myeloma proteins having

different binding sites 1041
BRACIALE, T. J. Immunologic recognition of influenza virus-infected cells. II. Expression of influenza A matrix protein on the infected cell surface and its role in recognition by cross-reactive cytotoxic T cells 673
BRADE, VOLKER, HALL, ROBERT E., and COLTEN, HARVEY R. Biosynthesis of pro-C3, a precursor of the third component of complement 759
BRAUN, DIETMAR G. See WEISS, KELUS, and BRAUN 1195
BROOKS, SUZANNE B. See KUNG, BROOKS, JAKWAY, LEONARD, and TALMAGE 665
BROWN, NORA M. See KIBLER, FRITZ, CHOU, CHOU, PEACOCKE, BROWN, and MCFARLIN 1323
BROXMEYER, HAL E. See RALPH, BROXMEYER, and NAKOINZ 611
BUCHHAGEN, DOROTHY L. See HANAFUSA, HALPERN, BUCHHAGEN, and KAWAI 1735
BURAKOFF, STEVEN. See BILLINGS, BURAKOFF, DORF, and BENACERRAF 623

C

CALLERY, RICHARD T. See STRELKAUSKAS, WILSON, CALLERY, CHESS, and SCHLOSSMAN 1765
CALNE, R. Y. See HOBART, LACHMAN, and CALNE 629
CAMBIER, J. C., VITETTA, E. S., KETTMAN, J. R., WETZEL, G. M., and UHR, J. W. B-cell tolerance. III. Effect of papain-mediated cleavage of cell surface IgD on tolerance susceptibility of murine B cells 107
—. See VITETTA, CAMBIER, LIGLER, KETTMAN, and UHR 1804
CAMPBELL, P. S. See MOSIER, MATHIESON, and CAMPBELL 59
CAREY, KATHLEEN. See HALL, COLVIN, CAREY, and MCCLUSKEY 1246
CAZENAVE, P. A. See BONA and CAZENAVE 881
CEBRA, JOHN. See OZATO, CEBRA, and EBERT 779
CHAKRAVARTY, ASHIM K., and CLARK, WILLIAM R. Lectin-driven maturation of cytotoxic effector cells: the nature of effector memory 230
CHESS, LEONARD. See STRELKAUSKAS, WILSON, CALLERY, CHESS, and SCHLOSSMAN 1765
—. See THORLEY-LAWSON, CHESS, and STROMINGER 495
CHIEN, C. C. See LIEBERMAN, VRANA, HUMPHREY, CHIEN, and POTTER 1294
CHIORAZZI, NICHOLAS, TUNG, AMAR S., and KATZ, DAVID H. Induction of a ragweed-specific allergic state in *Ir*-gene-restricted nonresponder mice 302
CHOU, FRANK C.-H. See KIBLER, FRITZ, CHOU, CHOU, PEACOCKE, BROWN, and MCFARLIN 1323
—. See WHITAKER, CHOU, CHOU, and KIBLER 317
CHOU, C.-H. JEN. See KIBLER, FRITZ, CHOU, CHOU, PEACOCKE, BROWN, and MCFARLIN 1323
—. See WHITAKER, CHOU, CHOU, and KIBLER 317
CHU, LILY. See KASTNER, RICH, CHU, and RICH 1152
CLAGETT, JAMES. See PRESS, ROSSE, and CLAGETT 735
CLAMAN, HENRY N., MILLER, STEPHEN D., and SY, MAN-SUN. Suppressor cells in tolerance to contact sensitivity active against hapten-syngeneic and hapten-allogeneic determinants 49
CLARK, WILLIAM R. See CHAKRAVARTY and CLARK 230
CLINE, MARTIN J., and BILLING, RONALD. Antigens expressed by human B lymphocytes and myeloid stem cells 1143
COHEN, JOEL E., D'EUSTACHIO, PETER, and EDELMAN, GERALD M. The specific antigen-binding cell populations of individual fetal mouse spleens: repertoire composition, size, and genetic control 394
COHN, ZANVIL A. See JOHNSON, MEI, and COHN 1613
—. See NOGUEIRA, GORDON, and COHN 157
—. See NOGUEIRA, GORDON, and COHN 172
COLOMBANI, M. J. See GOMARD, DUPREZ, REME, COLOMBANI, and LEVY 909
COLTEN, HARVEY R. See BRADE, HALL, and COLTEN 759
COLVIN, ROBERT B. See HALL, COLVIN, CAREY, and MCCLUSKEY 1246
COMINGS, DAVID E. See RUOSLAHTI, ENGVALL, JALANKO, and COMINGS 1054
CONE, ROBERT E., GERSHON, RICHARD K., and ASKENASE, PHILIP W. Nylon adherent antigen-specific rosette-forming T cells 1390
COOPER, MAX D. See KEARNEY, COOPER, KLEIN, ABNEY, PARKHOUSE, and LAWTON 297
—. See MORETTA, WEBB, GROSSI, LYDYARD, and COOPER 184
COUTINHO, ANTONIO. See EICHMANN, COUTINHO, and MELCHERS 1436
CRADDOCK, JOHN G. See CROMARTIE, CRADDOCK, SCHWAB, ANDERLE, and YANG 1585
CRAMER, MATTHIAS. See KRAWINKEL, CRAMER, MAGE, KELUS, and RAJEWSKY 792
CROMARTIE, WILLIAM J., CRADDOCK, JOHN G., SCHWAB, JOHN H., ANDERLE, SONIA K., and YANG, CHANG-HSU. Arthritis in rats after systemic injection of streptococcal cells or cell walls 1585
CUMMINGS, K. B. See BEAN, KODERA, CUMMINGS, and BLOOM 1455

D

DANNEMAN, PAMELA JONES, and MICHAEL, J. GABRIEL. Regulation of reaginic antibody production in mice. I. Suppression by antigen of IgE antibody production in vitro 1534

DAVID, CHELLA S. See OKUDA, DAVID, and SHREFFLER 1561
—. See PONZIO, DAVID, SHREFFLER, and THORBECKE 132
DAVIE, JOSEPH M. See WARREN and DAVIE 1627
DEAK, BEVERLY. See MERUELO, DEAK, and MCDEVITT 1367
—. See MERUELO, LIEBERMAN, DEAK, and MCDEVITT 1088
—. See MERUELO, LIEBERMAN, GINZTON, DEAK, and MCDEVITT 1079
DELEO, ALBERT B., SHIKU, HIROSHI, TAKAHASHI, TOSHITADA, JOHN, MARY, and OLD, LLOYD J. Cell surface antigens of chemically induced sarcomas of the mouse. I. Murine leukemia virus-related antigens and alloantigens on cultured fibroblasts and sarcoma cells: description of a unique antigen on BALB/c Meth A sarcoma 720
DELOVITCH, TERRY L., and MCDEVITT, HUGH O. In vitro analysis of allogeneic lymphocyte interaction. I. Characterization and cellular origin of an Ia-positive helper factor-allogeneic effect factor 1019
—, MURPHY, DONAL B., and MCDEVITT, HUGH O. Immunochemical evidence for three Ia loci in the *I*-region of the *H-2* complex 1549
D'EUSTACHIO, PETER. See COHEN, D'EUSTACHIO, and EDELMAN 394
DE WITT, C. See BOSMA, DE WITT, HAUSMAN, MARKS, and POTTER 1041
DICKLER, HOWARD B., AHMED, AFTAB, and SACHS, DAVID H. B-lymphocyte Fc receptor-associated non-H-2 antigens are determined by a single polymorphic locus which is linked to the *Mls* locus 1678
—. See SINGER, DICKLER, and HODES 1096
DIERICH, MANFRED P., and LANDEN, BEATE. Complement bridges between cells. Analysis of a possible cell-cell interaction mechanism 1484
DOE, WILLIAM F. See RUSSELL, DOE, and MCINTOSH 1511
DOHERTY, PETER C. See BIDDISON, DOHERTY, and WEBSTER 690
—. See BIDDISON, SNODGRASS, BENNINK, EFFROS, and DOHERTY 617
—. See MARSHAK, DOHERTY, and WILSON 1773
DORF, MARTIN E., and STIMPFLING, JACK H. Coupled complementation of immune response genes controlling responsiveness to the H-2.2 alloantigen 571
—. See BILLINGS, BURAKOFF, DORF, and BENACERRAF 623
—. See GREENE, PIERRES, DORF, and BENACERRAF 293
—. See THÈZE, WALTENBAUGH, DORF, and BENACERRAF 287
DUBIN, THEODORE. See BECKER and DUBIN 457
DUPREZ, V. See GOMARD, DUPREZ, REME, COLOMBANI, and LEVY 909
DUROCHER, JOHN R. See MILLER, HAYNES, MCAULIFFE, SHIROISHI, DUROCHER, and MCGINNISS 277
DUTTON, RICHARD W. See TSE and DUTTON 747
DYMINSKI, J. W., and SMITH, R. T. Generation of cytotoxic effector cells by immunocompetent thymus-cell subpopulations 412

E

EBERT, JAMES D. See OZATO, CEBRA, and EBERT 779
EDELMAN, GERALD M. See COHEN, D'EUSTACHIO, and EDELMAN 394
—. See WEBB, GALL, and EDELMAN 923
EFFROS, RITA B. See BIDDISON, SNODGRASS, BENNINK, EFFROS, and DOHERTY 617
EICHMANN, KLAUS, COUTINHO, ANTONIO, and MELCHERS, FRITZ. Absolute frequencies of lipopolysaccharide-reactive B cells producing A5A idiotype in unprimed, streptococcal A carbohydrate-primed, anti-A5A idiotype-sensitized and anti-A5A idiotype-suppressed A/J mice 1436
EIG, BLAIR M., JU, SHYR-TE, and NISONOFF, ALFRED. Complete inhibition of the expression of an idiotype by a mechanism of B-cell dominance 1574
ENGVALL, EVA. See RUOSLAHTI, ENGVALL, JALANKO, and COMINGS 1054
ENNIS, FRANCIS A., MARTIN, W. JOHN, and VERBONITZ, MARTHA W. Hemagglutinin-specific cytotoxic T-cell response during influenza infection 893

F

FEARON, DOUGLAS T., and AUSTEN, K. FRANK. Activation of the alternative complement pathway with rabbit erythrocytes by circumvention of the regulatory action of endogenous control proteins 22
FEHR, JORG, and JACOB, HARRY S. In vitro granulocyte adherence and in vivo marginations: two associated complement-dependent functions. Studies based on the acute neutropenia of filtration leukophoresis 641
FENYÖ, E. M., YEFENOF, E., KLEIN, E., and KLEIN, G. Immunization of mice with syngeneic Moloney lymphoma cells induces separate antibodies against virion envelope glyco-protein and virus-induced cell surface antigens 1521
FERNANDEZ, CARMEN, and MÖLLER, GÖRAN. Induction of immunological tolerance requires that the B cells can respond to the polyclonal B-cell-activating properties of the thymus-independent antigens 308
—, and —. Immunological unresponsiveness to thymus-independent antigens: two fundamentally different genetic mechanisms of B-cell unresponsiveness to dextran 1663
FERREIRA, ARTURO, TAKAHASHI, MORINOBU,

and NUSSENZWEIG, VICTOR. Purification and characterization of mouse serum protein with specific binding affinity for C4 (Ss protein) 1001
FIEDLER, J. See PRANGE, FIEDLER, NITECKI, and BELLONE 766
FIELDS, BERNARD N. See WEINER and FIELDS 1305
FISCHETTI, VINCENT A. Streptococcal M protein extracted by nonionic detergent. II. Analysis of the antibody response to the multiple antigenic determinants of the M-protein molecule 1108
FORMAN, JAMES. Cytotoxic T cells distinguish between trinitrophenyl- and dinitrophenyl-modified syngeneic cells 600
FOX, EUGENE N. See YANG, SOPREY, WITTNER, and FOX 344
FRELINGER, JEFFREY A. See WETTSTEIN and FRELINGER 1356
—. See WETTSTEIN, HAUGHTON, and FRELINGER 1346
FRITZ, ROBERT B. See KIBLER, FRITZ, CHOU, CHOU, PEACOCKE, BROWN, and MCFARLIN 1323
FUDENBERG, H. HUGH. See GALBRAITH, GOUST, and FUDENBERG 1821
FUKS, ZVI. See SLAVIN, STROBER, FUKS, and KAPLAN 34

G

GALANOS, CHRIS. See HOFFMANN, GALANOS, KOENIG, and OETTGEN 1640
GALBRAITH, ROBERT M., GOUST, JEAN-MICHEL, and FUDENBERG, H. HUGH. Lymphocyte culture: induction of colonies by conditioned medium from human lymphoid cell lines 1821
GALL, W. EINAR. See WEBB, GALL, and EDELMAN 923
GEARHART, PATRICIA J. See SIGAL, PICKARD, METCALF, GEARHART, and KLINMAN 933
GECZY, ANDREW F. See SCHWARTZ, KASK, PAUL, GECZY, and SHEVACH 547
—. See SHEVACH, LUNDQUIST, GECZY, and SCHWARTZ 561
GEDDE-DAHL, T., JR. See TEISBERG, OLAISEN, JONASSEN, GEDDE-DAHL, and THORSBY 1380
GENOVESI, EUGENE V., MARX, PRESTON A., and WHEELOCK, E. FREDERICK. Antigenic modulation of Friend virus erythroleukemic cells in vitro by serum from mice with dormant erythroleukemia 520
GERMAIN, RONALD N. See PIERCE, GERMAIN, KAPP, and BENACERRAF 1827
GERSHON, RICHARD K. See CONE, GERSHON, and ASKENASE 1390
GILLIS, STEVEN, and SMITH, KENDALL A. In vitro generation of tumor-specific cytotoxic lymphocytes. Secondary allogeneic mixed tumor lymphocyte culture of normal murine spleen cells 468
GINZTON, NANCY. See MERUELO, LIEBERMAN, GINZTON, DEAK, and MCDEVITT 1079
GOLDSTEIN, IRA M., LIND, STUART, HOFFSTEIN, SYLVIA, and WEISSMANN, GERALD. Influence of local anesthetics upon human polymorphonuclear leukocyte function in vitro. Reduction of lysosomal enzyme release and superoxide anion production 483
GOMARD, E., DUPREZ, V., REME, T., COLOMBANI, M. J., and LEVY, J. P. Exclusive involvement of *H-2D*^{b} or *H-2K*^{d} product in the interaction between T-killer lymphocytes and syngeneic *H-2*^{b} or *H-2*^{d} viral lymphomas 909
GOODMAN, DEWITT S. See MORSE, WITTE, and GOODMAN 1791
GOODMAN, JOEL W. See LEWIS and GOODMAN 1
GOODWIN, JAMES S., BANKHURST, ARTHUR D., and MESSNER, RONALD P. Suppression of human T-cell mitogenesis by prostaglandin: existence of a prostaglandin-producing suppressor cell 1719
GORDON, ROBERT D., SAMELSON, LAWRENCE E., and SIMPSON, ELIZABETH. Selective response to H-Y antigen by F_1 female mice sensitized to F_1 male cells 606
GORDON, SIAMON. See NOGUEIRA, GORDON, and COHN 157
—. See NOGUEIRA, GORDON, and COHN 172
GOTSCHLICH, E. C. See SALIT and GOTSCHLICH 1169
—. See SALIT and GOTSCHLICH 1182
GOUST, JEAN-MICHEL. See GALBRAITH, GOUST, and FUDENBERG 1821
GRANELLI-PIPERNO, ANGELA, VASSALLI, JEAN-DOMINIQUE, and REICH, E. Secretion of plasminogen activator by human polymorphonuclear leukocytes. Modulation by glucocorticoids and other effectors 1693
GREENE, MARK I., PIERRES, ANNE, DORF, MARTIN E., and BENACERRAF, BARUJ. The *I-J* subregion codes for determinants on suppressor factor(s) which limit the contact sensitivity response to picryl chloride 293
GROSSI, CARLO E. See MORETTA, WEBB, GROSSI, LYDYARD, and COOPER 184

H

HALL, CLIVE L., COLVIN, ROBERT B., CAREY, KATHLEEN, and MCCLUSKEY, ROBERT T. Passive transfer of autoimmune disease with isologous IgG_1 and IgG_2 antibodies to the tubular basement membrane in strain XIII guinea pigs. Loss of self-tolerance induced by autoantibodies 1246
HALL, ROBERT E. See BRADE, HALL, and COLTEN 759
HALPER, J. P. See VANDE STOUWE, KUNKEL, HALPER, and WEKSLER 1809
HALPERN, CAROL C. See HANAFUSA, HALPERN, BUCHHAGEN, and KAWAI 1735
HALSTEAD, S. B., and O'ROURKE, E. J. Dengue viruses and mononuclear phagocytes. I. Infection enhancement by non-neutralizing antibody 201

HALSTEAD, S. B., O'ROURKE, E. J., and ALLISON, A. C. Dengue viruses and mononuclear phagocytes. II. Identity of blood and tissue leukocytes supporting in vitro infection 218

HAMAOKA, TOSHIYUKI, YOSHIZAWA, MUNETOSHI, YAMAMOTO, HIROSHI, KUROKI, MOTOMU, and KITAGAWA, MASAYASU. Regulatory functions of hapten-reactive helper and suppressor T lymphocytes. II. Selective inactivation of hapten-reactive suppressor T cells by hapten-nonimmunogenic copolymers of D-amino acids, and its application to the study of suppressor T-cell effect on helper T-cell development 91

—. See YAMAMOTO, HAMAOKA, YOSHIZAWA, KUROKI, and KITAGAWA 74

HAMMARSTRÖM, LENNART. See BICK, PERSSON, SMITH, MÖLLER, and HAMMARSTRÖM 1146

HANAFUSA, HIDESABURO, HALPERN, CAROL C., BUCHHAGEN, DOROTHY L., and KAWAI, SADAAKI. Recovery of avian sarcoma virus from tumors induced by transformation-defective mutants 1735

HARAN-GHERA, NECHAMA. See LONAI and HARAN-GHERA 1164

HARPEL, PETER C. Plasmin inhibitor interactions. The effectiveness of α_2-plasmin inhibitor in the presence of α_2-macroglobulin 1033

HASPEL, MARTIN V., PELLEGRINO, MICHELE A., LAMPERT, PETER W., and OLDSTONE, MICHAEL B. A. Human histocompatibility determinants and virus antigens: effect of measles virus infection on HLA expression 146

HAUGHTON, GEOFFREY. See WETTSTEIN, HAUGHTON, and FRELINGER 1346

HAUSMAN, S. J. See BOSMA, DE WITT, HAUSMAN, MARKS, and POTTER 1041

HAYNES, J. DAVID. See MILLER, HAYNES, MCAULIFFE, SHIROISHI, DUROCHER, and MCGINNISS 277

HEDRICK, STEPHEN. See MOCHIZUKI, HEDRICK, WATSON, and KINGSBURY 1500

HENKART, PIERRE A., SCHMITT-VERHULST, ANNE-MARIE, and SHEARER, GENE M. Specificity of cytotoxic effector cells directed against trinitrobenzene sulfonate-modified syngeneic cells. Failure to recognize cell surface-bound trinitrophenyl dextran 1068

HESTON, W. E., and PARKS, W. P. Mammary tumors and mammary tumor virus expression in hybrid mice of strains C57BL and GR 1206

HOBART, M. J., LACHMANN, P. J., and CALNE, R. Y. C6: synthesis by the liver in vivo 629

HODES, RICHARD J. See SINGER, DICKLER, and HODES 1096

HOFFMANN, MICHAEL K., GALANOS, CHRIS, KOENIG, SCOTT, and OETTGEN, HERBERT F. B-cell activation by lipopolysaccharide. Distinct pathways for induction of mitosis and antibody production 1640

HOFFSTEIN, SYLVIA. See GOLDSTEIN, LIND, HOFFSTEIN, and WEISSMANN 483

HUMPHREY, GLEN W. See SACHS, HUMPHREY, and LUNNEY 381

HUMPHREY, W., JR. See LIEBERMAN, VRANA, HUMPHREY, CHIEN, and POTTER 1294

I

IHLE, J. N. See MCCLINTOCK, IHLE, and JOSEPH 422

INMAN, JOHN K. See MOND, LIEBERMAN, INMAN, MOSIER, and PAUL 1138

J

JACEWICZ, MARY. See KEUSCH and JACEWICZ 535

JACOB, HARRY S. See FEHR and JACOB 641

JAKWAY, JAMES P. See KUNG, BROOKS, JAKWAY, LEONARD, and TALMAGE 665

JALANKO, HANNU. See RUOSLAHTI, ENGVALL, JALANKO, and COMINGS 1054

JOHANSSON, BJØRN. See WESTERMARK, NATVIG, and JOHANSSON 631

JOHN, MARY. See DELEO, SHIKU, TAKAHASHI, JOHN, and OLD 720

JOHNSON, WARREN D., JR., MEI, BARBARA, and COHN, ZANVIL A. The separation, long-term cultivation, and maturation of the human monocyte 1613

JONASSEN, R. See TEISBERG, OLAISEN, JONASSEN, GEDDE-DAHL, and THORSBY 1380

JONES, PATRICIA P. Analysis of H-2 and Ia molecules by two-dimensional gel electrophoresis 1261

JOSEPH, D. R. See MCCLINTOCK, IHLE, and JOSEPH 422

JU, SHYR-TE. See EIG, JU, and NISONOFF 1574

K

KANG, ANDREW H. See TRENTHAM, TOWNES, and KANG 857

KAPLAN, ALAN M., and MOHANAKUMAR, T. Expression of a new cell surface antigen on activated murine macrophages 1461

KAPLAN, HENRY S. See SLAVIN, STROBER, FUKS, and KAPLAN 34

KAPP, JUDITH A. See PIERCE, GERMAIN, KAPP, and BENACERRAF 1827

—. See WALTENBAUGH, THÈZE, KAPP, and BENACERRAF 970

KAPPAS, ATTALLAH. See MAINES and KAPPAS 1286

KAPPLER, JOHN W., and MARRACK, PHILIPPA. The role of *H-2*-linked genes in helper T-cell function. I. In vitro expression in B cells of immune response genes controlling helper T-cell activity 1748

KAROL, ROBIN A., REICHLIN, MORRIS, and NOBLE, ROBERT W. Evolution of an idiotypic determinant: anti-Val 435

KASK, ANNE M. See SCHWARTZ, KASK, PAUL, GECZY, and SHEVACH 547

KASTNER, DANIEL L., RICH, ROBERT R., CHU, LILY, and RICH, SUSAN SOLLIDAY. Regulatory mechanisms in cell-mediated immune re-

sponses. V. *H-2* homology requirements for the production of a minor locus-induced suppressor factor 1152
KATZ, DAVID H. See CHIORAZZI, TUNG, and KATZ 302
KAWAI, SADAAKI. See HANAFUSA, HALPERN, BUCHHAGEN, and KAWAI 1735
KEARNEY, J. F., COOPER, M. D., KLEIN, J., ABNEY, E. R., PARKHOUSE, R. M. E., and LAWTON, A. R. Ontogeny of Ia and IgD on IgM-bearing B lymphocytes in mice 297
KEES, URSULA. See BLANDEN, MCKENZIE, KEES, MELVOLD, and KOHN 869
KELUS, ANDREW S. See KRAWINKEL, CRAMER, MAGE, KELUS, and RAJEWSKY 792
—. See WEISS, KELUS, and BRAUN 1195
KETTMAN, J. R. See CAMBIER, VITETTA, KETTMAN, WETZEL, and UHR 107
—. See VITETTA, CAMBIER, LIGLER, KETTMAN, and UHR 1804
KEUSCH, GERALD T., and JACEWICZ, MARY. Pathogenesis of Shigella diarrhea. VII. Evidence for a cell membrane toxin receptor involving $\beta 1 \rightarrow 4$-linked *N*-acetyl-D-glucosamine oligomers 535
KIBLER, ROBERT F., FRITZ, ROBERT B., CHOU, FRANK C.-H., CHOU, C.-H. JEN, PEACOCKE, NANCILLE Y., BROWN, NORA M., and MCFARLIN, DALE E. Immune response of Lewis rats to peptide C1 (residues 68–88) of guinea pig and rat myelin basic proteins 1323
—. See WHITAKER, CHOU, CHOU, and KIBLER 317
KIKUTANI, HITOSHI. See NISHIZAWA, KISHIMOTO, KIKUTANI, and YAMAMURA 653
KINCADE, P. W. See KURLAND, KINCADE, and MOORE 1420
KINGSBURY, DAVID T. See MOCHIZUKI, HEDRICK, WATSON, and KINGSBURY 1500
KISHIMOTO, TADAMITSU. See NISHIZAWA, KISHIMOTO, KIKUTANI, and YAMAMURA 653
KITAGAWA, MASAYASU. See HAMAOKA, YOSHIZAWA, YAMAMOTO, KUROKI, and KITAGAWA 91
—. See YAMAMOTO, HAMAOKA, YOSHIZAWA, KUROKI, and KITAGAWA 74
KLEIN, E. See FENYÖ, YEFENOF, KLEIN, and KLEIN 1521
KLEIN, G. See FENYÖ, YEFENOF, KLEIN, and KLEIN 1521
KLEIN, J. See KEARNEY, COOPER, KLEIN, ABNEY, PARKHOUSE, and LAWTON 297
—. See PIERCE and KLINMAN 509
—. See SIGAL, PICKARD, METCALF, GEARHART, and KLINMAN 933
KOCH, CAMERON J. See MACDONALD and KOCH 698
KODERA, Y. See BEAN, KODERA, CUMMINGS, and BLOOM 1455
KOENIG, SCOTT. See HOFFMANN, GALANOS, KOENIG, and OETTGEN 1640
KOHN, HENRY I. See BLANDEN, MCKENZIE, KEES, MELVOLD, and KOHN 869
KOZAK, ROBERT. See WEKSLER and KOZAK 1833
KRAWINKEL, ULRICH, CRAMER, MATTHIAS, MAGE, ROSE G., KELUS, ANDREW S., and RAJEWSKY, KLAUS. Isolated hapten-binding receptors of sensitized lymphocytes. II. Receptors from nylon wool-enriched rabbit T lymphocytes lack serological determinants of immunoglobulin constant domains but carry the *A* locus allotypic markers 792
KUHN, CHARLES III. See WHITE, LIN, and KUHN 802
KUNG, JOHN T., BROOKS, SUZANNE B., JAKWAY, JAMES P., LEONARD, LUISE L., and TALMAGE, DAVID W. Suppression of in vitro cytotoxic response by macrophages due to induced arginase 665
KUNKEL, H. G. See VANDE STOUWE, KUNKEL, HALPER, and WEKSLER 1809
KURLAND, J. I., KINCADE, P. W., and MOORE, M. A. S. Regulation of B-lymphocyte clonal proliferation by stimulatory and inhibitory macrophage-derived factors 1420
KUROKI, MOTOMU. See HAMAOKA, YOSHIZAWA, YAMAMOTO, KUROKI, and KITAGAWA 91
—. See YAMAMOTO, HAMAOKA, YOSHIZAWA, KUROKI, and KITAGAWA 74
KWA, B. H., and LIEW, F. Y. Immunity in taeniasis-cysticercosis. I. Vaccination against *Taenia taeniaeformis* in rats using purified antigen 118

L

LACHMANN, P. J. See HOBART, LACHMANN, and CALNE 629
LAMM, MICHAEL E. See ROUX, MCWILLIAMS, PHILLIPS-QUAGLIATA, WEISZ-CARRINGTON, and LAMM 1311
LAMPERT, PETER W. See HASPEL, PELLEGRINO, LAMPERT, and OLDSTONE 146
LANDEN, BEATE. See DIERICH and LANDEN 1484
LAWTON, A. R. See KEARNEY, COOPER, KLEIN, ABNEY, PARKHOUSE, and LAWTON 297
LAYTON, JUDITH E. See SCOTT, LAYTON, and NOSSAL 1473
LEHTO, V.-P. See LINDER, LEHTO, VIRTANEN, STENMAN, and NATVIG 1158
LEONARD, LOUISE L. See KUNG, BROOKS, JAKWAY, LEONARD, and TALMAGE 665
LEVY, J. P. See GOMARD, DUPREZ, REME, COLOMBANI, and LEVY 909
LEWIS, GEORGE K., and GOODMAN, JOEL W. Carrier-directed anti-hapten responses by B-cell subsets 1
LIEBERMAN, MIRIAM. See MERUELO, LIEBERMAN, DEAK, and MCDEVITT 1088
—. See MERUELO, LIEBERMAN, GINZTON, DEAK, and MCDEVITT 1079
LIEBERMAN, ROSE, VRANA, M., HUMPHREY, W. JR., CHIEN, C. C., and POTTER, M. Idiotypes of inulin-binding myeloma proteins localized to variable region light and heavy chains: genetic significance 1294

LIEBERMAN, ROSE, VRANA, M., HUMPHREY, W. JR., CHIEN, C. C., and POTTER, M. See MOND, LIEBERMAN, INMAN, MOSIER, and PAUL 1138
LIEW, F. Y. See KWA and LIEW 118
LIGLER, F. S. See VITETTA, CAMBIER, LIGLER, KETTMAN, and UHR 1804
LIN, HSIU-SAN. See WHITE, LIN, and KUHN 802
LIND, STUART. See GOLDSTEIN, LIND, HOFFSTEIN, and WEISSMANN 483
LINDAHL, KIRSTEN FISCHER. See WILSON, LINDAHL, WILSON, and SPRENT 361
LINDER, E., LEHTO, V.-P., VIRTANEN, I., STENMAN, S., and NATVIG, J. B. Localization of amyloid-related serum protein SAA-like material to intermediate (10 nm) filaments of cultured human embryonal fibroblasts 1158
LONAI, PETER, and HARAN-GHERA, NECHAMA. Resistance genes to murine leukemia in the *I* immune response gene region of the *H-2* complex 1164
LUNDQUIST, MARILYN L. See SHEVACH, LUNDQUIST, GECZY, and SCHWARTZ 561
LUNNEY, JOAN K. See SACHS, HUMPHREY, and LUNNEY 381
LYDYARD, PETER M. See MORETTA, WEBB, GROSSI, LYDARD, and COOPER 184

M

MACDONALD, H. ROBSON. Energy metabolism and T-cell-mediated cytolysis. II. Selective inhibition of cytolysis by 2-deoxy-D-glucose 710
– and KOCH, CAMERON J. Energy metabolism and T-cell-mediated cytolysis. I. Synergism between inhibitors of respiration and glycolysis 698
MAGE, ROSE G. See KRAWINKEL, CRAMER, MAGE, KELUS, and RAJEWSKY 792
MAINES, MAHIN D., and KAPPAS, ATTALLAH. Enzymes of heme metabolism in the kidney. Regulation by trace metals which do not form heme complexes 1286
MARKOWETZ, B. See VAN SNICK, MARKOWETZ, and MASSON 817
MARKS, R. See BOSMA, DE WITT, HAUSMAN, MARKS, and POTTER 1041
MARRACK, PHILIPPA. See KAPPLER and MARRACK 1748
MARSHAK, ANN, DOHERTY, PETER C., and WILSON, DARCY B. The control of specificity of cytotoxic T lymphocytes by the major histocompatibility complex (AG-B) in rats and identification of a new alloantigen system showing no AG-B restriction 1773
MARTIN, W. JOHN. See ENNIS, MARTIN, and VERBONITZ 893
MARX, PRESTON A. See GENOVESI, MARX, and WHEELOCK 520
MASSON, P. L. See VAN SNICK, MARKOWETZ, and MASSON 817
MATHIESON, B. J. See MOSIER, MATHIESON, and CAMPBELL 59
MCAULIFFE, FLORENCE M. See MILLER, HAYNES, MCAULIFFE, SHIROISHI, DUROCHER, and MCGINNISS 277
MCCARTY, M. See VAN DE RIJN, ZABRISKIE, and MCCARTY 579
MCCLINTOCK, P. R., IHLE, J. N., and JOSEPH, D. R. Expression of AKR murine leukemia virus gp71-like and BALB(X) gp71-like antigens in normal mouse tissues in the absence of overt virus expression 422
MCCLUSKEY, ROBERT T. See HALL, COLVIN, CAREY, and MCCLUSKEY 1246
MCDEVITT, HUGH O. See DELOVITCH and MCDEVITT 1019
–. See DELOVITCH, MURPHY, and MCDEVITT 1549
–. See MERUELO, DEAK, and MCDEVITT 1367
–. See MERUELO, LIEBERMAN, DEAK, and MCDEVITT 1088
–. See MERUELO, LIEBERMAN, GINZTON, DEAK, and MCDEVITT 1079
–. See PRESS and MCDEVITT 1815
MCFARLIN, DALE E. See KIBLER, FRITZ, CHOU, CHOU, PEACOCKE, BROWN, and MCFARLIN 1323
MCGINNISS, MARY H. See MILLER, HAYNES, MCAULIFFE, SHIROISHI, DUROCHER, and MCGINNISS 277
MCINTOSH, ANN TOZIER. See RUSSELL, DOE, and MCINTOSH 1511
MCKENZIE, IAN F. C. See BLANDEN, MCKENZIE, KESS, MELVOLD, and KOHN 869
–. See PARISH and MCKENZIE 332
MCMICHAEL, A. J., and SASAZUKI, T. A suppressor T cell in the human mixed lymphocyte reaction 368
MCWILLIAMS, MICHAEL. See ROUX, MCWILLIAMS, PHILLIPS-QUAGLIATA, WEISZ-CARRINGTON, and LAMM 1311
MEI, BARBARA. See JOHNSON, MEI, and COHN 1613
MEINKE, G. C. See WALKER, MEINKE, and WEIGLE 445
MELCHERS, FRITZ. See EICHMANN, COUTINHO, and MELCHERS 1436
MELVOLD, ROGER W. See BLANDEN, MCKENZIE, KEES, MELVOLD, and KOHN 869
MERUELO, DANIEL, LIEBERMAN, MIRIAM, GINZTON, NANCY, DEAK, BEVERLY, and MCDEVITT, HUGH O. Genetic control of radiation leukemia virus-induced tumorigenesis. I. Role of the major murine histocompatibility complex, *H-2* 1079
–, –, DEAK, BEVERLY, and MCDEVITT, HUGH O. Genetic control of radiation leukemia virus-induced tumorigenesis. II. Influence of *Srlv-1*, a locus not linked to *H-2* 1088
–, DEAK, BEVERLY, and MCDEVITT, HUGH O. Genetic control of cell-mediated responsiveness to an AKR tumor-associated antigen. Mapping of the locus involved to the *I* region of the *H-2* complex 1367

MESSNER, RONALD P. See GOODWIN, BANKHURST, and MESSNER 1719
METCALF, ELEANOR S. See SIGAL, PICKARD, METCALF, GEARHART, and KLINMAN 933
MICHAEL, J. GABRIEL. See DANNEMAN and MICHAEL 1534
MILLER, J. F. A. P. See TANIGUCHI and MILLER 1450
MILLER, LOUIS H., HAYNES, J. DAVID, MCAULIFFE, FLORENCE M., SHIROISHI, TSUGIYE, DUROCHER, JOHN R., and MCGINNISS, MARY H. Evidence for differences in erythrocyte surface receptors for the malarial parasites, *Plasmodium falciparum* and *Plasmodium knowlesi* 277
MILLER, R. G. See TEH, PHILLIPS, and MILLER 1280
MILLER, STEPHEN D., See CLAMAN, MILLER, and SY 49
MOCHIZUKI, DIANE, HEDRICK, STEPHEN, WATSON, JAMES, and KINGSBURY, DAVID T. The interaction of herpes simplex virus with murine lymphocytes. I. Mitogenic properties of herpes simplex virus 1500
MOHANAKUMAR, T. See KAPLAN and MOHANAKUMAR 1461
MÖLLER, ERNA. See BICK, PERSSON, SMITH, MÖLLER, and HAMMARSTRÖM 1146
MÖLLER, GÖRAN. See BICK and MÖLLER 844
—. See FERNANDEZ and MÖLLER 308
—. See FERNANDEZ and MÖLLER 1663
MOND, JAMES J., LIEBERMAN, ROSE, INMAN, JOHN K., MOSIER, DONALD E., and PAUL, WILLIAM E. Inability of mice with a defect in B-lymphocyte maturation to respond to phosphorylcholine on immunogenic carriers 1138
MOORE, M. A. S. See KURLAND, KINCADE, and MOORE 1420
MORETTA, LORENZO, WEBB, SUSAN R., GROSSI, CARLO E., LYDYARD, PETER M., and COOPER, MAX D. Functional analysis of two human T-cell subpopulations: help and suppression of B-cell responses by T cells bearing receptors for IgM or IgG 184
MORSE, JANE H., WITTE, LARRY D., and GOODMAN, DEWITT S. Inhibition of lymphocyte proliferation stimulated by lectins and allogeneic cells by normal plasma lipoproteins 1791
MOSIER, D. E., MATHIESON, B. J., and CAMPBELL, P. S. Ly phenotype and mechanism of action of mouse neonatal suppressor T cells 59
—. See MOND, LIEBERMAN, INMAN, MOSIER, and PAUL 1138
—. See ZITRON, MOSIER, and PAUL 1707
MÜLLER-EBERHARD, HANS J. See PANGBURN, SCHREIBER, and MÜLLER-EBERHARD 257
MURPHY, DONAL B. See DELOVITCH, MURPHY, and MCDEVITT 1549

N

NAKOINZ, ILONA. See RALPH, BROXMEYER, and NAKOINZ 611
NATHAN, CARL F., and ROOT, RICHARD K. Hydrogen peroxide release from mouse peritoneal macrophages: dependence on sequential activation and triggering 1648
NATVIG, JACOB B. See LINDER, LEHTO, VIRTANEN, STENMAN, and NATVIG 1158
—. See WESTERMARK, NATVIG, and JOHANSSON 631
NISHIZAWA, YOSHIO, KISHIMOTO, TADAMITSU, KIKUTANI, HITOSHI, and YAMAMURA, YUICHI. Induction and properties of cytoplasmic factor(s) which enhance nuclear nonhistone protein phosphorylation in lymphocytes stimulated by anti-Ig 653
NISONOFF, ALFRED. See EIG, JU, and NISONOFF 1574
NITECKI, D. E. See PRANGE, FIEDLER, NITECKI, and BELLONE 766
NOBLE, ROBERT W. See KAROL, REICHLIN, and NOBLE 435
NOGUEIRA, NADIA, GORDON, SIAMON, and COHN, ZANVIL. *Trypanosoma cruzi:* modification of macrophage function during infection 157
—, GORDON, SIAMON, and COHN, ZANVIL. *Trypanosoma cruzi:* the immunological induction of macrophage plasminogen activator requires thymus-derived lymphocytes 172
NOSSAL, G. J. V. See SCOTT, LAYTON, and NOSSAL 1473
NUSSENZWEIG, VICTOR. See FERREIRA, TAKAHASHI, and NUSSENZWEIG 1001

O

OETTGEN, HERBERT F. See HOFFMANN, GALANOS, KOENIG, and OETTGEN 1640
OKUDA, KENJI, DAVID, CHELLA S., and SHREFFLER, DONALD C. The role of gene products of the *I-J* subregion in mixed lymphocyte reactions 1561
OKUMURA, KO, TAKEMORI, TOSHITADA, TOKUHISA, TAKESHI, and TADA, TOMIO. Specific enrichment of the suppressor T cell bearing I-J determinants. Parallel functional and serological characterizations 1234
OLAISEN, B. See TEISBERG, OLAISEN, JONASSEN, GEDDE-DAHL, and THORSBY 1380
OLD, LLOYD J. See DELEO, SHIKU, TAKAHASHI, JOHN, and OLD 720
—. See STOCKERT and OLD 271
OLDSTONE, MICHAEL B. A. See HASPEL, PELLEGRINO, LAMPERT, and OLDSTONE 146
—. See PERRIN, ZINKERNAGEL, and OLDSTONE 949
O'ROURKE, E. J. See HALSTEAD and O'ROURKE 201
—. See —, —, and ALLISON 218
ORSON, FRANK M. See RICH, ORSON, and RICH 1221
OZATO, KEIKO, CEBRA, JOHN, and EBERT, JAMES D. Cell-mediated mitogenic response induced by leukoagglutinin and *Lens culinaris* lectin in mouse lymphocytes 779

P

PANGBURN, MICHAEL K., SCHREIBER, ROBERT D., and MÜLLER-EBERHARD, HANS J. Human complement C3b inactivator: isolation, characterization, and demonstration of an absolute requirement for the serum protein β1H for cleavage of C3b and C4b in solution 257
PARISH, CHRISTOPHER R., and MCKENZIE, IAN F. C. Direct visualization of T lymphocytes bearing Ia antigens controlled by the *I-J* subregion 332
PARKHOUSE R. M. E. See KEARNEY, COOPER, KLEIN, ABNEY, PARKHOUSE, and LAWTON 297
PARKS, W. P. See HESTON and PARKS 1206
PAUL, WILLIAM E. See MOND, LIEBERMAN, INMAN, MOSIER, and PAUL 1138
—. See SCHWARTZ, KASK, PAUL, GECZY, and SHEVACH 547
—. See YANO, SCHWARTZ, and PAUL 828
—. See ZITRON, MOSIER, and PAUL 1707
PEACOCKE, NANCILLE Y. See KIBLER, FRITZ, CHOU, CHOU, PEACOCKE, BROWN, and MCFARLIN 1323
PELLEGRINO, MICHELE A. See HASPEL, PELLEGRINO, LAMPERT, and OLDSTONE 146
PERRIN, LUC H., ZINKERNAGEL, ROLF M., and OLDSTONE, MICHAEL B. A. Immune response in humans after vaccination with vaccinia virus: generation of a virus-specific cytotoxic activity by human peripheral lymphocytes 949
PERSSON, ULLA. See BICK, PERSSON, SMITH, MÖLLER, and HAMMARSTRÖM 1146
PHILLIPS, R. A. See TEH, PHILLIPS, and MILLER 1280
PHILLIPS-QUAGLIATA, JULIA M. See ROUX, MCWILLIAMS, PHILLIPS-QUAGLIATA, WEISZ-CARRINGTON, and LAMM 1311
PICKARD, ALLAN R. See SIGAL, PICKARD, METCALF, GEARHART, and KLINMAN 933
PIERCE, CARL W., GERMAIN, RONALD N., KAPP, JUDITH A., and BENACERRAF, BARUJ. Secondary antibody responses in vitro to L-glutamic acid^{60}-L-alanine^{30}-L-tyrosine^{10}(GAT) by (responder × nonresponder) F_1 spleen cells stimulated by parental GAT-macrophages 1827
PIERCE, SUSAN K., and KLINMAN, NORMAN R. Antibody-specific immunoregulation 509
PIERRES, ANNE, See GREENE, PIERRES, DORF, and BENACERRAF 293
PILARSKI, LINDA M. Ontogeny of cell-mediated immunity. I. Early development of alloantigen-specific cytotoxic T-cell precursors in postnatal mice 887
PISETSKY, DAVID S., and SACHS, DAVID H. The genetic control of the immune response to staphylococcal nuclease. VI. Recombination between genes determining the A/J anti-nuclease idiotypes and the heavy chain allotype locus 1603
PONZIO, N. M., DAVID, C. S., SHREFFLER, D. C., and THORBECKE, G. J. Properties of reticulum cell sarcomas in SJL/J mice. V. Nature of reticulum cell sarcoma surface antigen which induces proliferation of normal SJL/J T cells 132
POTTER, M. See BOSMA, DE WITT, HAUSMAN, MARKS, and POTTER 1041
—. See LIEBERMAN, VRANA, HUMPHREY, CHIEN, and POTTER 1294
POWELL, A. E. See SMITH and POWELL 899
PRANGE, C. A., FIEDLER, J., NITECKI, D. E., and BELLONE, C. J. Inhibition of T-antigen-binding cells by idiotypic antisera 766
PRESS, JOAN L., and MCDEVITT, HUGH O. Allotype-specific analysis of anti-(T,G)-A—L antibodies produced by *Ir-1A* high- and low-responder chimeric mice 1815
PRESS, OLIVER W., ROSSE, CORNELIUS, and CLAGETT, JAMES. Phytohemagglutinin-induced differentiation and blastogenesis of precursor T cells from mouse bone marrow 735

R

RAJEWSKY, KLAUS. See KRAWINKEL, CRAMER, MAGE, KELUS, and RAJEWSKY 792
RALPH, PETER, BROXMEYER, HAL E., and NAKOINZ, ILONA. Immunostimulators induce granulocyte/macrophage colony-stimulating activity and block proliferation in a monocyte tumor cell line 611
REICH, E. See GRANELLI-PIPERNO, VASSALLI, and REICH 1693
REICHLIN, MORRIS. See KAROL, REICHLIN, and NOBLE 435
REME, T. See GOMARD, DUPREZ, REME, COLOMBANI, and LEVY 909
RICH, ROBERT R. See KASTNER, RICH, CHU, and RICH 1152
—. See RICH, ORSON, and RICH 1221
RICH, SUSAN SOLLIDAY, ORSON, FRANK M., and RICH, ROBERT R. Regulatory mechanisms in cell-mediated immune responses. VI. Interaction of *H-2* and non-*H-2* genes in elaboration of mixed leukocyte reaction suppressor factor 1221
—. See KASTNER, RICH, CHU, and RICH 1152
ROOT, RICHARD K. See NATHAN and ROOT 1648
ROSSE, CORNELIUS. See PRESS, ROSSE, and CLAGETT 735
ROUX, MARIA E., MCWILLIAMS, MICHAEL, PHILLIPS-QUAGLIATA, JULIA M., WEISZ-CARRINGTON, PAUL, and LAMM, MICHAEL E. Origin of IgA-secreting plasma cells in the mammary gland 1311
RUOSLAHTI, ERKKI, ENGVALL, EVA, JALANKO, HANNU, and COMINGS, DAVID E. Antigenic differences in nuclear proteins of normal liver and hepatoma. Identification of a nuclear protein present in hepatocytes but absent in hepatoma cells 1054
RUSSELL, STEPHEN W., DOE, WILLIAM F., and MCINTOSH, ANN TOZIER. Functional characterization of a stable, noncytolytic stage of macrophage activation in tumors 1511

S

SACHS, DAVID H., HUMPHREY, GLEN W., and LUNNEY, JOAN K. Sharing of Ia antigens between species. I. Detection of Ia specificities shared by rats and mice 381
—. See DICKLER, AHMED, and SACHS 1678
—. See PISETSKY and SACHS 1603
SALIT, I. E., and GOTSCHLICH, E. C. Hemagglutination by purified type I *Escherichia coli* pili 1169
— and —. Type I *Escherichia coli* pili: characterization of binding to monkey kidney cells 1182
SAMELSON, LAWRENCE E. See GORDON, SAMELSON, and SIMPSON 606
SASAZUKI, T. See MCMICHAEL and SASAZUKI 368
SCHLOSSMAN, STUART F. See STRELKAUSKAS, WILSON, CALLERY, CHESS, and SCHLOSSMAN 1765
SCHMITT-VERHULST, ANNE-MARIE. See HENKART, SCHMITT-VERHULST, and SHEARER 1068
SCHREIBER, ROBERT D. See PANGBURN, SCHREIBER, and MÜLLER-EBERHARD 257
SCHWAB, JOHN H. See CROMARTIE, CRADDOCK, SCHWAB, ANDERLE, and YANG 1585
SCHWARTZ, BENJAMIN D., KASK, ANNE M., PAUL, WILLIAM E., GECZY, ANDREW F., and SHEVACH, ETHAN M. The guinea pig *I* region. I. A structural and genetic analysis 547
—. See SHEVACH, LUNDQUIST, GECZY, and SCHWARTZ 561
SCHWARTZ, RONALD H. See YANO, SCHWARTZ, and PAUL 828
SCOTT, DAVID W., LAYTON, JUDITH E., and NOSSAL, G. J. V. Role of IgD in the immune response and tolerance. I. Anti-δ pretreatment facilitates tolerance induction in adult B cells in vitro 1473
SENIK, ANNA, and BLOOM, BARRY R. Differentiation of memory T cells to virus plaque-forming cells and cytotoxic T lymphocytes 11
SHEARER, GENE M. See HENKART, SCHMITT-VERHULST, and SHEARER 1068
SHEVACH, ETHAN M., LUNDQUIST, MARILYN L., GECZY, ANDREW F., and SCHWARTZ, BENJAMIN D. The guinea pig *I* region. II. Functional analysis 561
—. See SCHWARTZ, KASK, PAUL, GECZY, and SHEVACH 547
SHIKU, HIROSHI. See DELEO, SHIKU, TAKAHASHI, JOHN, and OLD 720
SHIROISHI, TSUGIYE. See MILLER, HAYNES, MCAULIFFE, SHIROISHI, DUROCHER, and MCGINNISS 277
SHREFFLER, DONALD C. See OKUDA, DAVID, and SHREFFLER 1561
—. See PONZIO, DAVID, SHREFFLER, and THORBECKE 132
SIGAL, NOLAN H. Novel idiotypic and antigen-binding characteristics in two anti-dinitrophenyl monoclonal antibodies 282
SIGAL, NOLAN H., PICKARD, ALLAN R., METCALF, ELEANOR S., GEARHART, PATRICIA J., and KLINMAN, NORMAN R. Expression of phosphorylcholine-specific B cells during murine development 933
SIMPSON, ELIZABETH. See GORDON, SAMELSON, and SIMPSON 606
SINGER, ALFRED, DICKLER, HOWARD B., and HODES, RICHARD J. Cellular and genetic control of antibody responses in vitro. II. Ir gene control of primary IgM responses to trinitrophenyl conjugates of poly-L-(Tyr,Glu)-poly-D,L-Ala—poly-L-Lys and poly-L-(His, Glu)-poly-D,L-Ala—poly-L-Lys 1096
SLAVIN, SHIMON, STROBER, SAMUEL, FUKS, ZVI, and KAPLAN, HENRY S. Induction of specific tissue transplantation tolerance using fractionated total lymphoid irradiation in adult mice: long-term survival of allogeneic bone marrow and skin grafts 34
SMITH, EDVARD. See BICK, PERSSON, SMITH, MÖLLER, and HAMMARSTRÖM 1146
SMITH, KENDALL A. See GILLIS and SMITH 468
SMITH, R. N., and POWELL, A. E. The adoptive transfer of pregnancy-induced unresponsiveness to male skin grafts with thymus-dependent cells 899
SMITH, R. T. See DYMINSKI and SMITH 412
SNODGRASS, H. RALPH. See BIDDISON, SNODGRASS, BENNINK, EFFROS, and DOHERTY 617
SOPREY, PANDU R., See YANG, SOPREY, WITTNER, and FOX 344
SPRENT, JONATHAN. See WILSON, LINDAHL, WILSON, and SPRENT 361
STENMAN, S. See LINDER, LEHTO, VIRTANEN, STENMAN, and NATVIG 1158
STIMPFLING, JACK H. See DORF and STIMPFLING 571
STOCKERT, ELISABETH, and OLD, LLOYD J. Preleukemic expression of TL antigens in X-irradiated C57BL/6 mice 271
STREILEIN, J. WAYNE, and WIESNER, JAMES. Influence of splenectomy on first set rejection reactions of C57BL/6 females to male skin isografts 809
STRELKAUSKAS, ANTHONY J., WILSON, BARRY S., CALLERY, RICHARD T., CHESS, LEONARD, and SCHLOSSMAN, STUART F. T-cell regulation of human peripheral blood B-cell responsiveness 1765
STROBER, SAMUEL. See SLAVIN, STROBER, FUKS, and KAPLAN 34
STROMINGER, JACK L. See THORLEY-LAWSON, CHESS, and STROMINGER 495
SY, MAN-SUN. See CLAMAN, MILLER, and SY 49

T

TADA, TOMIO. See OKUMURA, TAKEMORI, TOKUHISA, and TADA 1234
TAKAHASHI, MORINOBU. See FERREIRA, TAKAHASHI, and NUSSENZWEIG 1001

TAKAHASHI, TOSHITADA. See DELEO, SHIKU, TAKAHASHI, JOHN, and OLD 720
TAKEMORI, TOSHITADA. See OKUMURA, TAKEMORI, TOKUHISA, and TADA 1234
TALMAGE, DAVID W. See KUNG, BROOKS, JAKWAY, LEONARD, and TALMAGE 665
TANIGUCHI, MASARU, and MILLER, J. F. A. P. Enrichment of specific suppressor T cells and characterization of their surface markers 1450
TARTAKOFF, ALAN M., and VASSALLI, PIERRE. Plasma cell immunoglobulin secretion. Arrest is accompanied by alterations of the Golgi complex 1332
TEH, H.-S., PHILLIPS, R. A., and MILLER, R. G. Quantitative studies on the precursors of cytotoxic lymphocytes. III. The lineage of memory cells 1280
TEISBERG, P., OLAISEN, B., JONASSEN, R., GEDDE-DAHL, T. JR., and THORSBY, E. The genetic polymorphism of the fourth component of human complement: methodological aspects and a presentation of linkage and association data relevant to its localization in the HLA region 1380
THÈZE, JACQUES, WALTENBAUGH, CARL, DORF, MARTIN E., and BENACERRAF, BARUJ. Immunosuppressive factor(s) specific for L-glutamic acid^{50}-L-tyrosine^{50} (GT). II. Presence of *I-J* determinants on the GT-suppressive factor 287
—. See WALTENBAUGH, THÈZE, KAPP, and BENACERRAF 970
THORBECKE, G. J. See PONZIO, DAVID, SHREFFLER, and THORBECKE 132
THORLEY-LAWSON, DAVID A., CHESS, LEONARD, and STROMINGER, JACK L. Suppression of in vitro Epstein-Barr virus infection. A new role for adult human T lymphocytes 495
THORSBY, E. See TEISBERG, OLAISEN, JONASSEN, GEDDE-DAHL, and THORSBY 1380
TOKUHISA, TAKESHI. See OKUMURA, TAKEMORI, TOKUHISA, and TADA 1234
TOWNES, ALEXANDER S. See TRENTHAM, TOWNES, and KANG 857
TRENTHAM, DAVID E., TOWNES, ALEXANDER S., and KANG, ANDREW H. Autoimmunity to type II collagen: an experimental model of arthritis 857
TSE, HARLEY Y., and DUTTON, RICHARD W. Separation of helper and suppressor T lymphocytes. II. Ly phenotypes and lack of DNA synthesis requirement for the generation of concanavalin A helper and suppressor cells 747
TUNG, AMAR S. See CHIORAZZI, TUNG, and KATZ 302

U

UHR, J. W. See CAMBIER, VITETTA, KETTMAN, WETZEL, and UHR 107
—. See VITETTA, CAMBIER, LIGLER, KETTMAN, and UHR 1804

V

VAN DE RIJN, I., ZABRISKIE, J. B., and MCCARTY, M. Group A streptococcal antigens cross-reactive with myocardium. Purification of heart-reactive antibody and isolation and characterization of the streptococcal antigen 579
VANDE STOUWE, R. A., KUNKEL, H. G., HALPER, J. P., and WEKSLER, M. E. Autologous mixed lymphocyte culture reactions and generation of cytotoxic T cells 1809
VAN SNICK, J. L., MARKOWETZ, B., and MASSON, P. L. The ingestion and digestion of human lactoferrin by mouse peritoneal macrophages and the transfer of its iron into ferritin 817
VASSALLI, JEAN-DOMINIQUE. See GRANELLI-PIPERNO, VASSALLI, and REICH 1693
VASSALLI, PIERRE. See TARTAKOFF and VASSALLI 1332
VERBONITZ, MARTHA W. See ENNIS, MARTIN, and VERBONITZ 893
VIRTANEN, I. See LINDER, LEHTO, VIRTANEN, STENMAN, and NATVIG 1158
VITETTA, E. S., CAMBIER, J. C., LIGLER, F. S., KETTMAN, J. R., and UHR, J. W. B-cell tolerance. IV. Differential role of surface IgM and IgD in determining tolerance susceptibility of murine B cells 1804
—. See CAMBIER, VITETTA, KETTMAN, WETZEL, and UHR 107
VRANA, M. See LIEBERMAN, VRANA, HUMPHREY, CHIEN, and POTTER 1294

W

WALKER, SHARYN M., MEINKE, G. C., and WEIGLE, WILLIAM O. Enrichment of antigen-specific B lymphocytes by the direct removal of B cells not bearing specificity for the antigen 445
WALTENBAUGH, CARL, THÈZE, JAQUES, KAPP, JUDITH A., and BENACERRAF, BARUJ. Immunosuppressive factor(s) specific for L-glutamic acid^{50}-L-tyrosine^{50} (GT). III. Generation of suppressor T cells by a suppressive extract derived from GT-primed lymphoid cells 970
—. See THÈZE, WALTENBAUGH, DORF, and BENACERRAF 287
WARREN, ROBERT W., and DAVIE, JOSEPH M. Antigen mediation of a late-acting suppressor T-cell activity 1627
WATSON, JAMES. See MOCHIZUKI, HEDRICK, WATSON, and KINGSBURY 1500
WEBB, CYNTHIA G., GALL, W. EINAR, and EDELMAN, GERALD M. Synthesis and distribution of H-2 antigens in preimplantation mouse embryos 923
WEBB, SUSAN R. See MORETTA, WEBB, GROSSI, LYDYARD, and COOPER 184
WEBSTER, ROBERT G. See BIDDISON, DOHERTY, and WEBSTER 690

WEIGLE, WILLIAM O. See BERMAN and WEIGLE 241
—. See WALKER, MEINKE, and WEIGLE 445
WEINER, HOWARD L., and FIELDS, BERNARD N. Neutralization of reovirus: the gene responsible for the neutralization antigen 1305
WEISER, W. Y., and BANG, F. B. Blocking of in vitro and in vivo susceptibility to mouse hepatitis virus 1467
WEISS, SIEGFRIED, KELUS, ANDREW S., and BRAUN, DIETMAR G. Antibody response to the streptococcal group A-variant polysaccharide in BASILEA rabbits lacking κ-polypeptide chains 1195
WEISSMANN, GERALD. See GOLDSTEIN, LIND, HOFFSTEIN, and WEISSMANN 483
WEISZ-CARRINGTON, PAUL. See ROUX, MCWILLIAMS, PHILLIPS-QUAGLIATA, WEISZ-CARRINGTON, and LAMM 1311
WEKSLER, MARC E., and KOZAK, ROBERT. Lymphocyte transformation induced by autologous cells. V. Generation of immunologic memory and specificity during the autologous mixed lymphocyte reaction 1833
—. See VANDE STOUWE, KUNKEL, HALPER, and WEKSLER 1809
WESTERMARK, PER, NATVIG, JACOB B., and JOHANSSON, BJØRN. Characterization of an amyloid fibril protein from senile cardiac amyloid 631
WETTSTEIN, PETER J., and FRELINGER, JEFFREY A. *H-2* effects on cell-cell interactions in the response to single non-H-2 alloantigens. II. *H-2-D* region control of H-7.1-specific stimulator function in mixed lymphocyte culture and susceptibility to lysis by H-7.1-specific cytotoxic cells 1356
—, HAUGHTON, GEOFFREY, and FRELINGER, JEFFREY. *H-2* effects on cell-cell interactions in the response to single non-H-2 alloantigens. I. Donor *H-2D* region control of H-7.1-immunogenicity and lack of restriction in vivo 1346
WETZEL, G. M. See CAMBIER, VITETTA, KETTMAN, WETZEL, and UHR 107
WHEELOCK, E. FREDERICK. See GENOVESI, MARX, and WHEELOCK 520
WHITAKER, JOHN N., CHOU, C.-H. JEN, CHOU, FRANK C.-H., and KIBLER, ROBERT F. Molecular internalization of a region of myelin basic protein 317
WHITE, RICHARD, LIN, HSIU-SAN, and KUHN, CHARLES III. Elastase secretion by peritoneal exudative and alveolar macrophages 802
WIESNER, JAMES. See STREILEIN and WIESNER 809
WIGHT, R. See ANDERSSON, AGUET, WIGHT, ANDERSSON, BINZ, and WIGZELL 1124
WIGZELL, H. See ANDERSSON, AGUET, WIGHT, ANDERSSON, BINZ, and WIGZELL 1124
WILSON, BARRY S. See STRELKAUSKAS, WILSON, CALLERY, CHESS, and SCHLOSSMAN 1765
WILSON, DARCY B., LINDAHL, KIRSTEN FISCHER, WILSON, DIANNE H., and SPRENT, JONATHAN. The generation of killer cells to trinitrophenyl-modified allogeneic targets by lymphocyte populations negatively selected to strong alloantigens 361
—. See MARSHAK, DOHERTY, and WILSON 1773
WILSON, DIANNE H. See WILSON, LINDAHL, WILSON, and SPRENT 361
WITTE, LARRY D. See MORSE, WITTE, and GOODMAN 1791
WITTNER, MASAKO K. See YANG, SOPREY, WITTNER, and FOX 344

Y

YAMAMOTO, HIROSHI, HAMAOKA, TOSHIYUKI, YOSHIZAWA, MUNETOSHI, KUROKI, MOTOMU, and KITAGAWA, MASAYASU. Regulatory functions of hapten-reactive helper and suppressor T lymphocytes. I. Detection and characterization of hapten-reactive suppressor T-cell activity in mice immunized with hapten-isologous protein conjugate 74
—. See HAMAOKA, YOSHIZAWA, YAMAMOTO, KUROKI, and KITAGAWA 91
YAMAMURA, YUICHI. See NISHIZAWA, KISHIMOTO, KIKUTANI, and YAMAMURA 653
YANG, CHANG-HSU. See CROMARTIE, CRADDOCK, SCHWAB, ANDERLE, and YANG 1585
YANG, LILY C., SOPREY, PANDU R., WITTNER, MASAKO K., and FOX, EUGENE N. Streptococcal-induced cell-mediated immune destruction of cardiac myofibers in vitro 344
YANO, AKIHIKO, SCHWARTZ, RONALD H., and PAUL, WILLIAM E. Antigen presentation in the murine T-lymphocyte proliferative response. I. Requirement for genetic identity at the major histocompatibility complex 828
YEFENOF, E. See FENYÖ, YEFENOF, KLEIN, and KLEIN 1521
YOSHIZAWA, MUNETOSHI. See HAMAOKA, YOSHIZAWA, YAMAMOTO, KUROKI, and KITAGAWA 91
—. See YAMAMOTO, HAMAOKA, YOSHIZAWA, KUROKI, and KITAGAWA 74
YURT, ROGER, and AUSTEN, K. FRANK. Preparative purification of the rat mast cell chymase: characterization and interaction with granule components 1405

Z

ZABRISKIE, J. B. See VAN DE RIJN, ZABRISKIE, and MCCARTY 579
ZINKERNAGEL, ROLF M. See PERRIN, ZINKERNAGEL, and OLDSTONE 949
ZITRON, IAN M., MOSIER, DONALD E., and PAUL, WILLIAM E. The role of surface IgD in the response to thymic-independent antigens 1707
ZWERNER, ROBERT K., BARSTAD, PAUL A., and ACTON, RONALD T. Isolation and characterization of murine cell surface components. I. Purification of milligram quantities of Thy-1.1 986

INDEX TO SUBJECTS

A

N-Acetyl-D-glucosamine oligomers, $\beta 1 \rightarrow 4$-linked, involved in cell membrane toxin receptor in *Shigella* diarrhea, 535

Acid, deoxyribonucleic, synthesis not required for activation of cytotoxic T cells by polyribonucleotides, 844

Acid^{50}, L-glutamic-, L-tyrosine^{50}, specific immunosuppressive factors, generation of suppressor T cells by extract from primed lymphoid cells, 970

Acid^{60}-, L-glutamic, L-alanine^{30}-L-tyrosine^{10}, secondary antibody responses in vitro by (responder × nonresponder) spleen cells stimulated by parental macrophages, 1827

Acids, D-amino, hapten-nonimmunogenic copolymers, inactivation of hapten-reactive suppressor T cells, and suppressor T-cell effect on helper T-cell development, 91

L-Alanine^{30}-L-tyrosine^{10}, L-glutamic acid^{60}-secondary antibody responses in vitro by (responder × nonresponder) spleen cells stimulated by parental macrophages, 1827

poly-D,L-Alanine—poly-L-Lys, poly-L-(His, Glu)-, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096

poly-D,L-Alanine—poly-L-Lys, poly-L-(Tyr, Glu)-, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096

poly-D,L-Alanine—poly-L-lysine, poly-L-(tyrosine, glutamine)-, antibodies produced by high and low responders, allotype-specific analysis, 1815

Allergic state, ragweed-specific, induction in *Ir*-gene-restricted nonresponders, 302

Alloantigen, H-2.2, immune response genes controlling responsiveness, coupled complementation, 571

– system showing no AG-B restriction, in control of specificity, by major histocompatibility complex (AG-B), in cytotoxic T lymphocytes, 1773

Alloantigens and antigens, leukemia virus-related, on cultured fibroblasts and sarcoma cells, unique antigen on BALB/c Meth A sarcoma, 720

–, negatively selecting lymphocytes that generate killer cells to trinitrophenyl-modified allogeneic targets, 361

Alloantigens, non-H-2, single, H-2 effects on cell-cell interactions in response, donor *H-2D* region control of H-7.1 immunogenicity, lack of restriction in vivo, 1346

–, non-H-2, single, H-2 effects on cell-cell interactions in response, *H-2 D* region control of H-7.1-specific stimulator function in mixed lymphocyte culture, susceptibility to lysis by H-7.1-specific cytotoxic cells, 1356

Alloantigen-specific cytotoxic T-cell precursors, early postnatal development, 887

Allogeneic and syngeneic, hapten-, determinants, contact sensitivity active against, suppressor cells in tolerance, 49

– bone marrow and skin grafts, long-term survival, induction of specific tissue transplantation tolerance using fractionated total lymphoid irradiation, 34

– cells and lectins, stimulation of lymphocyte proliferation inhibited by normal plasma lipoproteins, 1791

– effect factor, Ia-positive helper factor-, in allogeneic lymphocyte interaction in vitro, 1019

– lymphocyte interaction in vitro, Ia-positive helper factor-allogeneic effect factor, 1019

– mixed tumor lymphocyte culture, secondary, generating tumor-specific cytotoxic lymphocytes, 468

– targets, trinitrophenyl-modified, to killer cells generated by lymphocytes negatively selected to alloantigens, 361

Allotype-specific analysis of anti-poly-L-(tyrosine, glutamine)-poly-D,L-alanine—poly-L-lysine antibodies produced by high and low responders, 1815

Allotypic markers, *a* locus, carried by hapten-binding receptors of sensitized T lymphocytes, 792

– suppression, maternally induced, release by nocardia water-soluble mitogen, 881

Alveolar and peritoneal exudative macrophages, elastase secretion, 802

D-Amino acids, hapten-nonimmunogenic copolymers, inactivation of hapten-reactive suppressor T cells, and suppressor T-cell effect on helper T-cell development, 91

Amyloid, cardiac, senile, and amyloid fibril protein, 631

– fibril protein, from senile cardiac amyloid, 631

Amyloid-related serum protein SAA-like material, localization to intermediate filaments of cultured human embryonal fibroblasts, 1158

Anesthetics, local, influence upon human polymorphonuclear leukocyte function in vitro, reducing lysosomal enzyme release and superoxide anion production, 483

Antibodies, isologous, IgG_1 and IgG_2, to tubular basement membrane, and passive trans-

fer of autoimmune disease, 1246
Antibodies, monoclonal, anti-dinitrophenyl, idiotypic and antigen-binding characteristics, 282
–, separate, against virion envelope glycoprotein and virus-induced cell-surface antigens, induced by immunization with syngeneic Moloney lymphoma cells, 1521
– to poly-L-(tyrosine, glutamine)-poly-D,L-alanine–poly-L-lysine, produced by high and low responders, allotype-specific analysis 1815
Antibody, heart-reactive, and group A streptococcal antigen, cross-reactive with myocardium, 579
–, IgE, suppression of production by antigen, in vitro, 1534
–, influenza virus matrix protein, detecting common antigen on surface of cells infected with influenza A viruses, 690
–, non-neutralizing, enhancement of dengue virus infection in mononuclear phagocytes, 201
– production and mitosis, induction pathways, in B-cell activation by lipopolysaccharide, 1640
– response to multiple antigenic determinants of streptococcal M-protein molecule, 1108
– response to streptococcal group A-variant polysaccharide, in subjects lacking K-polypeptide chains, 1195
– responses in vitro, cellular and genetic control, 1096
– responses, secondary, in vitro, to L-glutamic acid^{60}-L-alanine^{30}-L-tyrosine^{10} by (responder × nonresponder) spleen cells stimulated by parental macrophages, 1827
Antibody-specific immunoregulation, 509
Antigen, AKR tumor-associated, cell-mediated responsiveness, genetic control, locus mapped to *I* region of *H-2* complex, 1367
–, cell surface, expression on activated macrophages, 1461
–, common, cell surface, in infection with influenza A viruses, detected by antibody to influenza virus matrix protein, 690
–, group A streptococcal, cross-reactive with myocardium, and heart-reactive antibody, 579
–, H-Y, selective response by F_1 females sensitized to F_1 male cells, 606
– mediation of late-acting suppressor T-cell activity, 1627
– neutralizing reovirus, gene responsible, 1305
– presentation in murine T-lymphocyte proliferative response, requiring genetic identity at major histocompatibility complex, 828
– suppression of IgE antibody production in vitro, 1534
–, surface, nature of, in reticulum cell sarcoma, inducing proliferation of normal T cells, 132
–, unique, on cells of BALB/c Meth A sarcoma, 720
Antigen-binding and idiotypic characteristics, in anti-dinitrophenyl monoclonal antibodies, 282
– cell populations of individual fetal spleens, 394
Antigen-, T, binding cells, inhibition by idiotypic antisera, 766
Antigenic determinants, multiple, of streptococcal M-protein molecule, antibody response, 1108
– differences between normal liver and hepatoma, nuclear protein present in hepatocytes, absent in hepatoma, 1054
– modulation of Friend virus erythroleukemic cells in vitro, by serum from mice with dormant erythroleukemia, 520
Antigens and alloantigens, leukemia virus-related, on cultured fibroblasts and sarcoma cells, unique antigen on BALB/c Meth A sarcoma, 720
–, cell-surface, virus-induced, and virion envelope glycoproteins, separate antibodies induced by immunization with syngeneic Moloney lymphoma cells, 1521
– expressed by B lymphocytes and myeloid stem cells, in man, 1143
–, gp71-like and BALB(X) gp71-like, to AKR murine leukemia virus, in normal tissues, in absence of overt virus expression, 422
–, group A streptococcal, cross-reactive with myocardium, purification of heart-reactive antibody, isolation of streptococcal antigen, 579
–, *H-2*, in preimplantation embryos, synthesis and distribution, 923
–, Ia, and IgD, ontogeny on IgM-bearing B lymphocytes, 297
–, Ia, controlled by *I-J* subregion, on T lymphocytes, 332
–, Ia, specificities shared by two species, 381
–, non-H-2, Fc receptor-associated, B lymphocyte, determined by single polymorphic locus linked to *Mls* locus, 1678
–, TL, preleukemic expression in X-irradiated mice, 271
–, thymic-independent, surface IgD in response, 1707
–, thymus-independent, immunological unresponsiveness, B-cell unresponsiveness to dextran, genetic mechanisms, 1663
–, thymus-independent, polyclonal B cell-activating properties, B-cell response and induction of immunological tolerance, 308
–, virus, and histocompatibility determinants in man, effect of measles virus infection on HLA expression, 146
Antigen-specific B lymphocytes enriched by direct removal of B cells not specific for antigen, 445
– T cells, rosette-forming, nylon-adherent, 1390
Anti-hapten responses by B-cell subsets, carrier-directed, 1

Antisera, idiotypic, and inhibition of T antigen-binding cells, 766
Arginase, induced, and suppression of in vitro cytotoxic response by macrophages, 665
Arthritis, after systemic injection of streptococcal cells or cell walls, 1585
–, experimental model, autoimmunity to type II collagen, 857
Autoimmune disease, passive transfer, with isologous IgG_1 and IgG_2 antibodies to tubular basement membrane, 1246
Autoimmunity to type II collagen, arthritis experimental model, 857
Autoimmunogen in inducing specific immune unresponsiveness, use of purified mixed leukocyte culture-activated T lymphoblasts, 1124

B

B-cell activation by lipopolysaccharide, pathways for induction of mitosis and antibody production, 1640
– dominance mechanism, expression of idiotype, complete inhibition, 1574
– response to polyclonal B-cell-activating properties of thymus-independent antigens, and induction of immunological tolerance, 308
– responses in man, help and suppression by T cells bearing receptors for IgM or IgG, functional analysis, 184
– subsets, carrier-directed anti-hapten responses, 1
– tolerance susceptibility, effect of papain-mediated cleavage of cell surface IgD, 107
– unresponsiveness, in human peripheral blood, T-cell regulation, 1765
– unresponsiveness to dextran, genetic mechanisms, 1663
B cells, adult, anti-δ pretreatment, facilitating tolerance induction in vitro, role of IgD in immune response, 1473
–, expressing *Ir* genes in vitro, 1748
–, lipopolysaccharide-reactive, producing A5A idiotype in unprimed, streptococcal A carbohydrate-primed, anti-A5A idiotype-sensitized, and anti-A5A idiotype-suppressed subjects, 1436
– not specific for antigen, direct removal enriching antigen-specific B lymphocytes, 445
–, phosphorylcholine-specific, expression during development, 933
–, tolerance susceptibility, and surface IgM and IgD, 1804
B-lymphocyte clonal proliferation, regulation by macrophage-derived factors, 1420
– Fc receptor-associated non-*H-2* antigens, determined by single polymorphic locus linked to *Mls* locus, 1678
– maturation defect, subjects nonresponsive to phosphorylcholine on immunogenic carriers, 1138
–, activation by Fc region of IgG, 241
B-lymphocytes, antigen-specific, enriched by direct removal of B cells not specific for antigen, 445
–, IgM-bearing, ontogeny of Ia antigens and IgD, 297
– and myeloid stem cells, expression of antigens, in man, 1143
Benzene sulfonate-, trinitro-, modified syngeneic cells, cytotoxic effector cells and failure to recognize cell surface-bound trinitrophenyl dextran, 1068
Blastogenesis and differentiation, phytohemagglutinin-induced, of precursor T cells, from bone marrow, 735
Blood, peripheral, human, T-cell regulation of B-cell unresponsiveness, 1765
– and tissue leukocytes, supporting in vitro infection of dengue virus in mononuclear phagocytes, identity, 218
Bone marrow, blastogenesis and differentiation of precursor T cells, phytohemagglutinin-induced, 735
– marrow and skin grafts, allogeneic, long-term survival, induction of specific tissue transplantation tolerance using fractionated total lymphoid irradiation, 34

C

Carbohydrate-(streptococcal A)-primed, unprimed, anti-A5A idiotype sensitized, anti-A5A idiotype-suppressed subjects, lipopolysaccharide-reactive B cells producing A5A idiotype, absolute frequencies, 1436
Cardiac myofibers in vitro, streptococcal-induced cell-mediated-immune destruction, 344
Cell-cell interaction, in complement bridges between cells, 1484
Cell surface component, Thy-1.1, purification of milligram quantities, 986
Chain, heavy, allotype locus, and anti-nuclease idiotypes, recombination between determining genes, 1693
Chains, heavy, of myeloma proteins, having different binding sites, common V_H determinants, 1041
–, light and heavy, variable region, localization of idiotypes of inulin-binding myeloma proteins, genetic significance, 1294
–, K-polypeptide, lack, and antibody response to streptococcal group A-variant polysaccharide, 1195
Choline, phosphoryl-, on immunogenic carriers, nonresponse by subjects of B-lymphocyte maturation defect, 1138
–, phosphoryl-, specific B cells, expression during development, 933
Chymase, mast cell, preparation, 1405
Clonal proliferation of B lymphocytes, regulation by macrophage-derived factors, 1420
Collagen, type II, autoimmunity, arthritis experimental model, 857
Complement bridges between cells, cell-cell interaction mechanism, 1484

Complement, human, fourth component, genetic polymorphism, and localization in HLA region, 1380
– inactivator in man, requirement for serum protein β1H for cleavage of C3b and C4b in solution, 257
– pathway, alternative, activation with erythrocytes by circumvention of regulatory action of endogenous control proteins, 22
–, precursor of third component, pro-C3, biosynthesis, 759
– 6, synthesis by liver in vivo, 629
Complementation, coupled, of immune response genes controlling responsiveness to H-2.2 alloantigen, 571
Concanavalin A helper and suppressor cells, Ly phenotypes and lack of DNA synthesis requirement for generation, separation of helper and suppressor T lymphocytes, 747
–, lack of response by lipopolysaccharide nonresponders, in genetic control of lymphocyte activation, 1146
Copolymers, hapten-nonimmunogenic, of D-amino acids, inactivation of hapten-reactive suppressor T cells, and suppressor T-cell effect on helper T-cell development, 91
Cysticercocis-, taeniasis, immunity, vaccination against *Taenia taeniformis* using purified antigen, 118
Cytolysis, T-cell-mediated, and energy metabolism, synergism between inhibitors of respiration and glycolysis, 698
–, T-cell-mediated, inhibition by 2-deoxy-D-glucose, 710
Cytotoxic activity, virus-specific, generated by human peripheral lymphocytes, in immune response to vaccinia virus vaccination in man, 949
– cells, H-7.1-specific, lysis, and H-7.1-specific stimulator function, in mixed lymphocyte culture, *H-2D* region control, 1356
– effector cells directed against trinitrobenzene sulfonate-modified syngeneic cells, failure to recognize cell surface-bound trinitrophenyl dextran, 1068
– effector cells generated by immunocompetent thymus cell subpopulations, 412
– effector cells, lectin-driven maturation, nature of effector memory, 230
– lymphocytes, precursors, lineage of memory cells, 1280
– lymphocytes, tumor-specific, generated by secondary allogeneic mixed tumor lymphocyte culture, 468
– response by macrophages, suppression in vitro due to induced arginase, 665
– T-cell activation by polyribonucleotides, DNA synthesis not required, 844
– T-cell precursors, alloantigen-specific, early postnatal development, 887
– T-cell response to ectromelia virus-infected cells, *H-2* requirement for precursor T-cell induction or lysis by effector T cells defined by BALB/*c-H-2*^{db} mutation, 869
Cytotoxic T-cell response, hemagglutinin-specific, in influenza, 893
– T cells, cross-reactive, and expression of influenza A matrix protein on cell surface, 673
– T cells, distinguishing between trinitrophenyl- and dinitrophenyl-modified syngeneic cells, 600
– T cells, generation, and autologous mixed lymphocyte culture reactions, 1809
– T cells, recognition of virus-specific modifications, induction not altered by substitution of target cells with trinitrophenol, 617
– T lymphocytes induced against allogeneic *I*-region determinants, reacting with Ia molecules on trinitrophenyl-conjugated syngeneic target cells, 623
– T lymphocytes, major histocompatibility complex (AG-B) controlling specificity, alloantigen system showing no AG-B restriction, 1773
– T lymphocytes and virus plaque-forming cells, differentiation of memory T cells, 11

D

Dengue viruses and mononuclear phagocytes, enhancement by non-neutralizing antibody, 201
– viruses and mononuclear phagocytes, identity of blood and tissue leukocytes supporting in vitro infection, 218
2-Deoxy-D-glucose, and inhibition of T-cell-mediated glycolysis, 710
Deoxyribonucleic acid, synthesis not required for activation of cytotoxic T cells by polyribonucleotides, 844
Dextran, B-cell unresponsiveness, genetic mechanisms, 1663
–, trinitrophenyl, cell surface-bound, not recognized by cytotoxic effector cells directed against trinitrobenzene sulfonate-modified syngeneic cells, 1068
Diarrhea, *Shigella*, pathogenesis, cell membrane toxin receptor involving β1 → 4-linked *N*-acetyl-D-glucosamine oligomers, 535
Dinitrophenyl, anti-, monoclonal antibodies, idiotypic and antigen-binding characteristics, 282

E

Ectromelia virus-infected cells, and cytotoxic T-cell response, *H-2* requirement for precursor T-cell induction or lysis by effector T cells defined by BALB/*c-H-2*^{db} mutation, 869
Elastase, secretion by peritoneal exudative and alveolar macrophages, 802
Electrophoresis, gel, two-dimensional, analysis of H-2 and Ia molecules, 1261
Embryonal fibroblasts, human, cultured, localization of amyloid-related serum protein SAA-like material to intermediate filaments, 1158
Embryos, preimplantation, and H-2 antigens, synthesis and distribution, 923

Enzymes of heme metabolism in kidney, regulation by trace metals not forming heme complexes, 1286
Epstein-Barr virus, suppression in vitro, role of adult human T lymphocytes, 495
Erythrocyte surface receptors for malarial parasites, *Plasmodium falciparum* and *Plasmodium knowlesi*, differences, 277
Erythrocytes activating alternative complement pathway by circumvention of regulatory action of endogenous control proteins, 22
Erythroleukemia, dormant, in mice, antigenic modulation by serum, in vitro, of Friend virus erythroleukemic cells, 520
Erythroleukemic cells, Friend virus, antigenic modulation in vitro by serum from mice with dormant erythroleukemia, 520
Escherichia coli, type I, pili, and binding to monkey kidney cells, 1182
–, type I, pili, and hemagglutination, 1169

F

Ferritin, and transfer of iron from human lactoferritin, 817
Fetal spleens, individual, specific antigen-binding cell populations, repertoire composition, size, genetic control, 394
Fibroblasts, human embryonal, cultured, intermediate filaments, and localization of amyloid-related serum protein SAA-like material, 1158
– and sarcoma cells, cultured, and leukemia virus-related antigens and alloantigens, unique antigen on BALB/c Meth A sarcoma, 720
Filtration leukophoresis leukopenia, acute, and granulocyte adherence in vitro, margination in vivo, 641
Friend virus erythroleukemic cells, antigenic modulation in vitro, by serum from mice with dormant erythroleukemia, 520

G

Gene, *Ir*-, restricted nonresponders, induction of ragweed-specific allergic state, 302
– products of *I-J* subregion, in mixed lymphocyte reactions, 1561
– region, *Ir*, of *H-2* complex, and leukemia resistance genes, 1164
Genes, determining A/J idiotypes and heavy chain allotype locus, recombination, in immune response to staphylococcal nuclease, 1603
–, *H-2* linked, in helper T-cell function, B cells expressing *Ir* genes in vitro, 1748
–, *H-2* and non-*H-2*, interacting in elaboration of mixed leukocyte reaction suppressor factor, in cell-mediated immune responses, 1221
–, immune response, controlling responsiveness to H-2.2 alloantigen, coupled complementation, 571
–, resistance to leukemia, in *Ir* gene region of *H-2* complex, 1164
Genetic and cellular control of antibody responses in vitro, 1096
– control of cell-mediated responsiveness to AKR tumor-associated antigen, mapping of locus involved to *I* region of *H-2* complex, 1367
– control of lymphocyte activation, nonresponse to concanavalin A in lipopolysaccharide nonresponders, 1146
– control of radiation leukemia virus-induced tumorigenesis, influence of *Srlv-1*, locus not linked to *H-2*, 1088
– control of radiation leukemia virus-induced tumorigenesis, role of major histocompatibility complex, *H-2*, 1079
– control, size, repertoire composition of specific antigen-binding cell populations, in individual fetal spleens, 394
– identity at major histocompatibility complex, required for antigen presentation in murine T-lymphocyte proliferative response, 828
– mechanisms of B-cell unresponsiveness to dextran, immunological unresponsiveness to thymus-independent antigens, 1663
– polymorphism of fourth component of human complement, localization in HLA region, 1380
– significance of localization, to variable region light and heavy chains, of idiotypes of inulin-binding myeloma proteins, 1294
– and structural analysis of guinea pig *I* region, 547
Glucocorticoids and other effectors, modulating plasminogen activator secretion by polymorphonuclear leukocytes, in man, 1791
D-Glucosamine, *N*-acetyl-, $\beta 1 \rightarrow 4$-linked, involved in cell membrane toxin receptor in *Shigella* diarrhea, 535
Glucose, 2-deoxy-D-, and inhibition of T-cell-mediated glycolyis, 710
L-Glutamic acid^{50}-L-tyrosine^{50}, presence of *I-J* determinants on specific immunosuppressive factor, 287
L-Glutamic acid^{50}-L-tyrosine^{50}, specific immunosuppressive factors, generation of suppresor T cells by extract from primed lymphoid cells, 970
L-Glutamic acid^{60}-L-alanine^{30}-L-tyrosine^{10}, secondary antibody responses in vitro by (responder × nonresponder) spleen cells stimulated by parental macrophages, 1827
Glutamine)-poly-D,L-alanine – poly-L-lysine, poly-L-(tyrosine,-, antibodies produced by high and low responders, allotype-specific analysis, 1815
Glutamine)-poly-D,L-Ala – poly-L-Lys, poly-L-(His,-, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
–)-poly-D,L-Ala – poly-L-Lys, poly-L-(tyr,-, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
Glycolysis and respiration, synergism between

inhibitors, energy metabolism and T-cell-mediated cytolysis, 698
Glycoprotein, tobacco, activating factor XII, 457
–, virion envelope, and virus-induced cell-surface antigens, separate antibodies induced by immunization with syngeneic Moloney lymphoma cells, 1521
Golgi complex, alterations accompanying arrest of plasma cell immunoglobulin secretion, 1332
Granulocyte/macrophage colony-stimulating activity induced by immunostimulators, 611
Granulocytes, in vitro adherence, in vivo margination, and acute neutropenia of filtration leukophoresis, 641

H

H-2 complex, *I* region, immunochemical evidence for three Ia loci, 1549
H-Y antigen, selective response by F_1 females sensitized to F_1 male cells, 606
Hapten, anti-, carrier-directed responses by B-cell subsets, 1
Hapten-binding receptors of sensitized T lymphocytes, lacking serological determinants of Ig constant domains, but carrying *a* locus allotypic markers, 792
Hapten-isologous protein conjugate, detection and characterization of hapten-reactive T-cell activity under immunization, 74
Hapten-nonimmunogenic copolymers of D-amino acids, inactivation of hapten-reactive suppressor T cells, suppressor T-cell effect on helper T-cell development, 91
Hapten-reactive suppressor T-cell activity, detection and characterization under immunization with hapten-isologous protein conjugate, 74
– suppressor T cells, inactivation by hapten-nonimmunogenic copolymers of D-amino acids, suppressor T-cell effect on helper T-cell development, 91
Hapten-syngeneic and allogeneic determinants, contact sensitivity active against, suppressor cells in tolerance, 49
Heart-reactive antibody and streptococcal antigen, group A streptococcal antigens cross-reactive with myocardium, 579
Hemagglutination by type I *E. coli* pili, 1169
Hemagglutinin-specific cytotoxic T-cell response in influenza, 893
Heme metabolism in kidney, enzymes, regulation by trace metals not forming heme complexes, 1286
Hepatitis virus, blocking of susceptibility in vivo and in vitro, 1467
Hepatoma and normal liver, antigenic differences, nuclear protein present in hepatocytes, absent in hepatoma, 1054
Herpes simplex virus, mitogenic properties, in interaction with lymphocytes, 1500
poly-L-(Histidine, Glu)-poly-D,L-Ala – poly-L-Lys, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
Histocompatibility complex, major, (AG-B), controling specificity of cytotoxic T lymphocytes, alloantigen system showing no AG-B restriction, 1773
– complex, major, genetic identity required for antigen presentation, in murine T-lymphocyte proliferative response, 828
– complex, major, *H-2*, role in genetic control of radiation leukemia virus-induced tumorigenesis, 1079
– determinants and virus antigens in man, effect of measles virus infection on HLA expression, 146
Histone, non-, nuclear, protein phosphorylation, enhancement in lymphocytes stimulated by anti-Ig, 653
Hydrogen peroxide release from peritoneal macrophages, sequential activation and triggering, 1735

I

Idiotype, A5A, produced by lipopolysaccharide-reactive B cells, in unprimed, streptococcal A carbohydrate-primed, anti-A5A idiotype-sensitized, and anti-A5A idiotype-suppressed subjects, 1436
–, expression by mechanism of B-cell dominance, complete inhibition, 1574
Idiotypes of inulin-binding myeloma proteins, localization to variable region light and heavy chains, 1294
Idiotypic and antigen-binding characteristics in anti-dinitrophenyl monoclonal antibodies, 282
– determinant, anti-Val, evolution, 435
Immune destruction, cell-mediated, streptococcal-induced, of cardiac myofibers in vitro, 344
– response, *Ir* gene control of primary IgM responses to trinitrophenyl conjugates of poly-L-(Tyr,Glu)-poly-D,L-Ala – poly-L-Lys and poly-L-(His,Glu)-poly-D,L-Ala – poly-L-Lys, 1096
– response, *Ir* gene region of *H-2* complex, and leukemia resistance genes, 1164
– response, *Ir* gene-restricted nonresponders, induction of ragweed-specific allergic state, 302
– response, *Ir* genes controlling responsiveness to H-2.2 alloantigen, coupled complementation, 571
– response, *Ir* genes, expression by B cells in vitro, 1748
– response and IgD, anti-δ pretreatment facilitating tolerance induction, in adult B cells in vitro, 1473
– response, *I*-region determinants, allogeneic, cytotoxic T lymphocytes react with Ia molecules on trinitrophenyl-conjugated syngeneic target cells, 623
– response, *I* region of guinea pig, functional analysis, 561

Immune response, *I* region of guinea pig, structural and genetic analysis, 547
Immune response, *I*-region of *H-2* complex, three Ia loci, immunological evidence, 1549
– response of Lewis rats to peptide C1 of guinea pig and rat myelin basic proteins, 1323
– response in man after vaccination with vaccinia virus, virus-specific cytotoxic activity generated by human peripheral lymphocytes, 949
– response to staphylococcal nuclease, recombination between genes determining A/J anti-nuclease idiotypes and heavy chain allotype locus, 1603
– responses, cell-mediated, *H-2* homology required for producing locus-induced suppressor factor, 1152
– responses, cell-mediated, *H-2* and non-*H-2* genes interacting in elaboration of mixed leukocyte reaction suppressor factor, 1221
– unresponsiveness, specific, induction using purified mixed leukocyte culture-activated T lymphoblasts as autoimmunogen, 1124
Immunity, auto-, to type II collagen, arthritis experimental model, 857
–, cell-mediated, early postnatal development of alloantigen-specific cytotoxic T cell precursors, 887
Immunization with syngeneic Moloney lymphoma cells, induction of separate antibodies against virion envelope glycoprotein and virus-induced cell-surface antigens, 1521
Immunochemical evidence for three Ia loci in *I* region of *H-2* complex, 1549
Immunocompetent thymus cell subpopulations generating cytotoxic effector cells, 412
Immunogenic carriers of phosphorylcholine, nonresponse by subjects of B-lymphocyte maturation defect, 1138
–, non-, hapten-, copolymers of D-amino acids, inactivation of hapten-reactive suppressor T cells, and suppressor T-cell effect on helper T cell development, 91
Immunogenicity, H-7.1-, donor *H-2D* region control, lack of restriction in vivo, 1346
Immunoglobulin A-secreting plasma cells in mammary gland, origin, 1311
–, anti-, stimulated lymphocytes, and enhancement of nuclear nonhistone protein phosphorylation, 653
– constant domains, serological determinants lacking in hapten-binding receptors of sensitized T lymphocytes, 792
– D, cell surface, effect of papain-mediated cleavage on tolerance susceptibility of B cells, 107
– D and Ia antigens, ontogeny on IgM-bearing B lymphocytes, 297
– D in immune response, anti-δ pretreatment facilitating tolerance induction, in adult B cells in vitro, 1473
– D, surface, in response to thymic-independent antigens, 1707
Immunoglobulin E antibody production, suppression by antigen in vitro, 1534
– G, activation of B lymphocytes by Fc region, 241
– G or IgM, T cells bearing receptors, help and suppression of B-cell responses in man, functional analysis, 184
– G_1 and G_2, isologous antibodies to tubular basement membrane, and passive transfer of autoimmune disease, 1246
– M and IgD, surface, and tolerance susceptibility of B cells, 1804
– M or IgG, T cells bearing receptors, help and suppression of B-cell responses in man, functional analysis, 184
– M-bearing B lymphocytes, ontogeny of Ia antigens and IgD, 297
– M responses, primary, to trinitrophenyl conjugates of poly-L-(Tyr, Glu)-poly-D,L-Ala–poly-L-Lys and poly-L-(His, Glu)-poly-D,L-Ala–poly-L-Lys, *Ir* gene control, 1096
– secretion by plasma cells, arrest accompanied by alterations of Golgi complex, 1332
Immunoglobulins, mouse, and serological distinction of heavy chain variable regions, common V_H determinants on heavy chains of mouse myeloma proteins with different binding sites, 1041
Immunologic memory and specificity, generated during autologous mixed lymphocyte reaction, 1833
– memory and specificity, generation during autologous mixed lymphocyte reaction, transformation of lymphocytes, 1809
Immunological tolerance and B-cell response to polyclonal B-cell-activating properties of thymus-independent antigens, 308
– unresponsiveness to thymus-independent antigens, B-cell unresponsiveness to dextran, genetic mechanisms, 1663
Immunoregulation, antibody-specific, 509
Immunostimulators induce granulocyte/macrophage colony-stimulating activity, block proliferaton in monocyte tumor cell line, 611
Immunosuppression of L-glutamic acid^{50}-L-tyrosine^{50}, presence of *I–J* determinants on specific suppressive factor, 287
Immunosuppressive factors specific for L-glutamic acid^{50}-L-tyrosine^{50}, generation of suppressor T cells by extract from primed lymphoid cells, 970
Influenza A matrix protein, expression on cell surface, recognition by cross-reactive cytotoxic T cells, 673
– A viruses, common antigen on surface of infected cells detected by antibody to influenza virus matrix protein, 690
–, hemagglutinin-specific cytotoxic T-cell response, 893
– virus matrix protein antibody, detecting common antigen on surface of cells infected with influenza A viruses, 690
Inhibitors, synergistic, of respiration and gly-

colysis, in energy metabolism and T-cell-mediated cytolysis, 698
Inulin-binding myeloma proteins, localization to variable region light and heavy chains, idiotypes, 1294
Iron, transfer into ferritin, ingestion and digestion of human lactoferritin by mouse peritoneal macrophages, 817
Irradiation, fractionated total lymphoid, induction of specific tissue transplantation tolerance, long-term survival of allogeneic bone marrow and skin grafts, 34
Isografts, male skin, and first set rejection reactions of C57BL/6 females, influence of splenectomy, 809

K

Kidney, enzymes of heme metabolism, regulation by trace metals not forming heme complexes,, 1286
Killer cells to trinitrophenyl-modified allogeneic targets, generated by lymphocytes negatively selected to alloantigens, 361

L

Lactoferrin, human, ingestion and digestion by mouse peritoneal macrophages, transfer of its iron into ferritin, 817
Lectin, *Lens culinaris*, and leukoagglutinin, induction of cell-mediated mitogenic response in lymphocytes, 779
Lectin-driven maturation of cytotoxic effector cells, nature of effector memory, 230
Lectins and allogeneic cells, stimulation of lymphocyte proliferation inhibited by normal plasma lipoproteins, 1791
Lens culinaris lectin and leukoagglutinin, induction of cell-mediated mitogenic response in lymphocytes, 779
Leukemia, radiation, virus-induced tumorigenesis, genetic control, influence of *Srlv-1*, locus not linked to *H-2*, 1088
–, radiation, virus-induced tumorigenesis, genetic control, role of major histocompatibility complex, *H-2*, 1079
–, resistance genes in *Ir* gene region of *H-2* complex, 1164
– virus, AKR murine, gp71-like and BALB(X) gp71-like antigens in normal tissues in absence of overt virus expression, 422
– virus-related antigens and alloantigens on cultured fibroblasts and sarcoma cells, unique antigen on BALB/c Meth A sarcoma, 720
Leukemic, pre-, expression of TL antigens in X-irradiated mice, 271
Leukoagglutinin and *Lens culinaris* lectin, induction of cell-mediated mitogenic response in lymphocytes, 779
Leukocyte reaction, mixed, elaboration of suppressor factor, *H-2* and non-*H-2* genes interacting, in cell-mediated immune responses, 1221
Leukocytes, of blood and tissue, supporting in vitro infection by dengue virus in mononuclear phagocytes, 218
–, human polymorphonuclear, influence of local anesthetics upon function in vitro, by reducing lysosomal enzyme release and superoxide anion production, 483
–, polymorphonuclear, in man, secretion of plasminogen activator modulated by glucocorticoids and other effectors, 1791
Leukophoresis, filtration, acute leukopenia, and granulocyte adherence in vitro, margination in vivo, 641
Lipopolysaccharide, activation of B cells, pathways for induction of mitosis and antibody production, 1640
– nonresponders, lack of response to concanavalin A, in genetic control of lymphocyte activation, 1146
Lipolysaccharide-reactive B cells producing A5A idiotype in unprimed, streptococcal A carbohydrate-primed, anti-A5A idiotype-sensitized, and anti-A5A idiotype-suppressed subjects, absolute frequencies, 1436
Lipoproteins, plasma, normal, inhibiting lymphocyte proliferation stimulated by lectins and allogeneic cells, 1791
Liver, normal, and hepatoma, antigenic differences in nuclear proteins, 1054
–, synthesis of C6 in vivo, 629
Locus-induced suppressor factor, in cell-mediated immune responses, *H-2* homology requirements, 1152
Lymphoblasts, T, purified mixed leukocyte culture-activated, use as autoimmunogen in inducing specific immune unresponsiveness, 1124
Lymphocyte activation, genetic control, nonresponse to concanavalin A in lipopolysaccharide nonresponders, 1146
–, B, antigens, Fc receptor-associated, non-H-2, determined by single polymorphic locus linked to *Mls* locus, 1678
–, B, clonal proliferation, regulation by macrophage-derived factors, 1420
–, B, maturation defect, subjects nonresponsive to phosphorylcholine on immunogenic carriers, 1138
– culture, induction, by conditioned medium, of colonies from human lymphoid cell lines, 1821
– culture, mixed, *H-2D* region control of H-7.1-specific stimulator function and lysis by H-7.1-specific cytoxic cells, 1356
– culture, mixed tumor, allogeneic, secondary, generating tumor-specific cytotoxic lymphocytes, 468
– culture reactions, mixed, autologous, and generation of cytotoxic T cells, 1809
– interaction, allogeneic, in vitro, Ia-positive helper factor-allogeneic effect factor, 1019
– proliferation stimulated by lectins and allogeneic cells, inhibition by normal plasma

lipoproteins, 1791
Lymphocyte reaction, autologous, mixed, and generation of immunologic memory and specificity, 1833
– reactions, mixed, and *I–J* subregion gene products, 1561
Lymphocyte, T, murine, proliferative response, antigen presentation requiring genetic identity at major histocompatibility complex, 828
Lymphocytes, anti-Ig stimulated, and enhancement of nuclear nonhistone protein phosphorylation, 653
–, B, activation by Fc region of IgG, 241
–, B, antigen-specific, enriched by direct removal of B cells not specific for antigen, 445
–, B, IgM-bearing, ontogeny of Ia antigens and IgD, 297
–, B, and myeloid stem cells, expression of antigens in man, 1143
–, cell-mediated mitogenic response induced by leukoagglutinin and *Lens culinaris* lectin, 779
–, cytotoxic, generated by secondary allogeneic mixed tumor lymphocyte culture, 468
–, cytotoxic, precursors, lineage of memory cells, 1280
–, and mitogenic properties of *Herpes simplex* virus, 1500
– negatively selected to alloantigens generating killer cells to trinitrophenyl-modified allogeneic targets, 361
–, peripheral, in man, generating virus-specific cytotoxic activity, in immune response after vaccination with vaccinia virus, 949
–, T, bearing Ia antigens controlled by *I–J* subregion, 332
–, T, cytotoxic, induced against allogeneic *I*-region determinants, reacting with Ia molecules on trinitrophenyl-conjugated syngeneic target cells, 623
–, T, cytotoxic, major histocompatibility complex (AG-B) controlling specificity, alloantigen system showing no AG-B restriction, 1773
–, T, cytotoxic, and virus plaque-forming cells, differentiation of memory T cells, 11
–, T, human, adult, role in suppressing Epstein-Barr virus in vitro, 495
–, T, killer, and syngeneic *H-2*^{b} or *H-2*^{d} viral lymphomas, involvement of *H-2*^{b} or *H-2K*^{d} product in interaction, 909
–, T, sensitized, hapten-binding receptors lacking serological determinants of Ig constant domains, but carrying *a* locus allotypic markers, 792
–, T, separation of helpers and suppressors, Ly phenotypes and lack of DNA synthesis requirement for generation of concanavalin A helper and suppressor cells, 747
–, thymus-derived, requirement for immunological induction of macrophage plasminogen activator in *Trypanosoma cruzi* infection, 172
Lymphoid cell lines, human, induction of colonies by conditioned medium, 1821
Lymphoid cells, GT-primed, generation of suppressor T cells by extract, 970
– irradiation, fractionated total, induction of specific tissue transplantation tolerance, long-term survival of allogeneic bone marrow and skin grafts, 34
Lymphoma, Moloney, syngeneic cells, immunization inducing separate antibodies to virion envelope glycoprotein and virus-induced cell-surface antigens, 1521
Lymphomas, viral, *H-2*^{b} or *H-2*^{d}, syngeneic, and killer T lymphocytes, involvement of *H-2*^{b} or *H-2K*^{d} product in interaction, 909
poly-L-Lysine, poly-L-(His, Glu)-poly-D,L-Ala–trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
poly-L-Lysine, poly-L-(tyrosine, glutamine)-poly-D,L-alanine–, antibodies produced by high and low responders, allotype-specific analysis, 1815

M

M protein molecule, streptococcal, antibody response to multiple antigenic determinants, 1108
α_2-Macroglobulin, and effectiveness of α_2-plasmin inhibitor, 1033
Macrophage activation in tumors, stable noncytolytic stage, 1511
– function, modification during infection with *Trypanosoma cruzi*, 157
– plasminogen activator in *Trypanosoma cruzi* infection, immunological induction requiring thymus-derived lymphocytes, 172
–-derived factors, regulating clonal proliferation of B lymphocytes, 1420
/granulocyte colony stimulating activity induced by immunostimulators, 611
Macrophages, activated, and expression of cell surface antigen, 1461
–, parental, stimulation of (responder × nonresponder) spleen cells, and secondary antibody response in vitro to L-glutamic acid^{60}-L-alanine^{30}-L-tyrosine^{10}, 1827
–, peritoneal exudative and alveolar, secretion of elastase, 802
–, peritoneal, mouse, ingestion and digestion of human lactoferritin, 817
–, peritoneal, releasing hydrogen peroxide, sequential activation and triggering, 1735
–, suppression of in vitro cytotoxic response due to induced arginase, 665
Malarial parasites, *Plasmodium falciparum* and *Plasmodium knowlesi*, differences in erythrocyte surface receptors, 277
Man, B lymphocytes and myeloid stem cells, expression of antigens, 1143
–, complement, fourth component, genetic

polymorphism, localization in HLA region, 1380
Man, complement inactivator, requirement for serum protein β1H for cleavage of C3b and C4b in solution, 257
–, embryonal fibroblasts, cultured, intermediate filaments, and localization of amyloid-related serum protein SAA-like material, 1158
–, help and suppression of B-cell responses by T cells bearing receptors for IgM or IgG, functional analysis, 184
–, histocompatibility determinants and virus antigens, effect of measles virus infection on HLA expression, 146
–, immune response after vaccination with vaccinia virus, virus-specific cytotoxic activity generated by human peripheral lymphocytes, 949
–, lactoferrin, ingestion and digestion by mouse peritoneal macrophages, transfer of its iron into ferritin, 817
–, lymphoid cell lines, induction of colonies by conditioned medium, 1821
–, monocyte, separation, long-term cultivation, maturation, 1613
–, mixed lymphocyte reaction, and suppressor T cell, 368
–, peripheral blood, B-cell unresponsiveness, T-cell regulation, 1765
–, peripheral lymphocytes generating virus-specific cytotoxic activity, in immune response after vaccination with vaccinia virus, 949
–, polymorphonuclear leukocyte function in vitro, influence of local anesthetics reducing lysosomal enzyme release and superoxide anion production, 483
–, polymorphonuclear leukocytes, secretion of plasminogen activator, modulation by glucocorticoids and other effectors, 1693
–, suppressor T-cell activity, 1455
–, T-cell mitogenesis, suppression by prostaglandin, and prostaglandin-producing suppressor cell, 1719
–, T lymphocytes, adult, role in suppressing Epstein-Barr virus in vitro, 495
Mast cell chymase, preparation, 1405
Measles virus infection, effect on HLA expression, 146
Membrane, basement, tubular, isologous IgG_1 and IgG_2 antibodies, and passive transfer of autoimmune disease, 1246
–, cell, toxin receptor involving β1 → 4-linked *N*-acetyl-D-glucosamine oligomers, in *Shigella* diarrhea, 535
Memory cells, lineage, precursors of cytotoxic lymphocytes, 1280
–, effector, nature in lectin-driven maturation of cytotoxic effector cells, 218
– T cells, differentiation to virus plaque-forming cells and cytotoxic T lymphocytes, 11
Mitogen, nocardia, water-soluble, and release from maternally induced allotypic suppression, 881
Mitogenesis, T-cell, in man, suppression by prostaglandin, and prostaglandin-producing suppressor cell, 1719
Mitogenic response, cell-mediated, induction by leukoagglutinin and *Lens culinaris* lectin in lymphocytes, 779
Mitosis, and antibody production, induction pathways, in B-cell activation by lipopolysaccharide, 1640
Mixed tumor lymphocyte culture, secondary, allogeneic, generating tumor-specific cytotoxic lymphocytes, 468
Monoclonal antibodies, anti-dinitrophenyl, idiotypic and antigen-binding characteristics, 282
Monocyte, human, separation, long-term cultivation, maturation, 1613
– tumor cell line, proliferation blocked by immunostimulators, 611
Mononuclear phagocytes and dengue viruses, enhancement by non-neutralizing antibody, 201
– phagocytes and dengue viruses, identity of blood and tissue leukocytes supporting infection in vitro, 218
Myelin basic protein, molecular internalization, 317
– basic proteins, guinea pig and rat, peptide C1, immune response of Lewis rats, 1323
Myeloid stem cells and B lymphocytes, expression of antigens, in man, 1143
Myeloma proteins, inulin-binding, localization to variable region light and heavy chains, idiotypes, 1294
– proteins with different binding sites, common V_H determinants on heavy chains, in serological distinction of heavy chain variable regions of mouse immunoglobulins, 1041
Myofibers, cardiac, in vitro, streptococcal-induced cell-mediated immune destruction, 344
Mutants, transformation-defective, induction of tumors, recovery of avian sarcoma virus, 1735

N

Neonatal suppressor T cells, Ly phenotype and mechanism of action, 59
Neutropenia, acute, of filtration leukophoresis, and granulocyte adherence in vitro, margination in vivo, 641
Nocardia water-soluble mitogen, and release from maternally induced allotypic suppression, 881
Nuclease, staphylococcal, immune response, recombination between genes determining A/J anti-nuclease idiotypes and heavy chain allotype locus, 1603
Nucleic acid, deoxyribo-, synthesis not required for activation of cytotoxic T cells by

polyribonucleotides, 844
Nylon-adherent T cells, antigen-specific, rosette-forming, 1390
Nylon wool-enriched T lymphocytes, receptors lacking serological determinants of Ig constant domains, but carrying *a* locus allotypic markers, 792

O

Oligomers, $\beta 1 \rightarrow 4$-linked *N*-acetyl-D-glucosamine, involved in cell membrane toxin receptor in *Shigella* diarrhea, 535
Oxyribonucleic acid, de-, synthesis not required for activation of cytotoxic T cells by polyribonucleotides, 844

P

Papain-mediated cleavage of cell surface IgD, effect on tolerance susceptibility of B cells, 107
Parasites, malarial, *Plasmodium falciparum* and *Plasmodium knowlesi*, differences in erythrocyte surface receptors, 277
Peptide C1 (residues 68–88) of myelin basic proteins, immune response of Lewis rats, 1323
Phagocytes, mononuclear, and dengue viruses, enhancement by non-neutralizing antibody, 201
–, mononuclear, and dengue viruses, identity of blood and tissue leukocytes supporting infection in vitro, 218
Phenol, trinitro-, substitution for target cells not altering induction of virus-specific modifications recognized by cytotoxic T cells, 617
Phenotype, Ly, and mechanism of action of neonatal suppressor T cells, 59
Phenyl, anti-dinitro-, monoclonal antibodies, idiotypic and antigen-binding characteristics, 282
Phenyl-, trinitro-, conjugated syngeneic target cells, Ia molecules, reaction of cytotoxic T lymphocytes induced against allogeneic *I*-region determinants, 623
Phenyl, trinitro-, conjugates of poly-L-(Tyr, Glu)-poly-D,L-Ala–poly-L-Lys and poly-L-(His, Glu)-poly-D,L-Ala–poly-L-Lys, primary IgM responses, *Ir* gene control, 1096
Phenyl-, trinitro- and dinitro-, modified syngeneic cells, distinguished by cytotoxic T cells, 600
Phenyl-, trinitro-, modified allogeneic targets to killer cells generated by lymphocytes negatively selected to alloantigens, 361
Phosphorylation of nuclear nonhistone proteins, enhanced in anti-Ig-stimulated lymphocytes, 653
Phosphorylcholine on immunogenic carriers, nonresponse by subjects of B-lymphocyte maturation defect, 1138
Phosphorylcholine-specific B cells, expression during development, 933
Phytohemagglutinin-induced differentiation and blastogenesis of precursor T cells from bone marrow, 735
Picryl chloride, contact sensitivity response, *I–J* subregion codes for determinants on suppressor factors, 293
Pili, of *Escherichia coli* type I, binding to monkey kidney cells, 1182
–, of *Escherichia coli* type I, hemagglutination, 1169
Plaque-forming cells, virus, and cytotoxic T lymphocytes, differentiation of memory T cells, 11
Plasma cell immunoglobulin secretion, arrest accompanied by alterations of Golgi complex, 1332
– cells, IgA-secreting, in mammary gland, origin, 1311
–, normal, lipoproteins, inhibition of lymphocyte proliferation stimulated by lectins and allogeneic cells, 1791
α_2-Plasmin inhibitor, effectiveness in presence of α_2-macroglobulin, 1033
Plasminogen activator, macrophage, in *Trypanosoma cruzi* infection, immunological induction requiring thymus-derived lymphocytes, 172
– activator, secretion by human polymorphonuclear leukocytes, modulation by glucocorticoids, other effectors, 1791
Plasmodium falciparum and *Plasmodium knowlesi*, malarial parasites, differences in erythrocyte surface receptors, 277
Plasmodium knowlesi and *Plasmodium falciparum*, malarial parasites, differences in erythrocyte surface receptors, 277
Polyclonal B cell-activating properties of thymus-independent antigens, B-cell response and induction of immunological tolerance, 308
Poly-L-lysine, poly-L-(His, Glu)-poly-D,L-Ala–, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
Poly-L-lysine, poly-L-(Tyr, Glu)-poly-D,L-Ala–, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
Polymorphism, genetic, of fourth component of human complement, and localization in HLA region, 1380
κ-Polypeptide chains, lack, and antibody response to streptococcal group A-variant polysaccharide, 1195
Polyribonucleotides, activation of cytotoxic T cell, DNA synthesis not required, 844
Polysaccharide, streptococcal group A-variant, antibody response in subjects lacking K-polypeptide chains, 1195
Pregnancy-induced unresponsiveness to male skin grafts, adoptive transfer with thymus-dependent cells, 899
Preleukemic expression of TL antigens in X-irradiated mice, 271
Prostaglandin, suppression of T-cell mitogene-

sis, and prostaglandin-producing suppressor cell, in man, 1719
Prostaglandin-producing suppressor cell, and prostaglandin suppression of T-cell mitogenesis, in man, 1719
Protein, amyloid fibril, from senile cardiac amyloid, 631
– conjugate, hapten-isologous, detection and characterization of hapten-reactive T-cell activity under immunization, 74
–, influenza virus matrix, antibody detecting common antigen on surface of cells infected with influenza A viruses, 690
–, M-, molecule, streptococcal, antibody response to multiple antigenic determinants, 1108
–, matrix, influenza A, expression on cell surface, and recognition by cross-reactive cytotoxic T cells, 673
–, mouse serum, with specific binding affinity for C4, 1001
–, myelin basic, molecular internalization, 317
–, nuclear, present in hepatocytes, absent in hepatoma, 1054
– phosphorylation, nonhistone, nuclear, enhancement in lymphocytes stimulated by anti-Ig, 653
–, serum, β1H, a requirement for cleavage of human complements C3b and C4b in solution, 257
–, serum, SAA-like material, amyloid-related, localization to intermediate filaments of cultured human embryonal fibroblasts, 1158
Proteins, basic, myelin, guinea pig and rat, peptide C1, immune response of Lewis rats, 1323
–, control, endogenous, activation of alternative complement pathway with rabbit erythrocytes by circumvention of regulatory action, 22
–, myeloma, inulin-binding, localization to variable region light and heavy chains, idiotypes, 1294

R

Radiation leukemia virus-induced tumorigenesis, genetic control, influence of *Srlv-1*, locus not linked to *H-2*, 1088
– leukemia virus tumorigenesis, genetic control, role of major histocompatibilty complex, *H-2*, 1079
Ragweed-specific allergic state, induction in *Ir* gene-restricted nonresponders, 302
Reaginic antibody production, suppression by antigen of IgE antibody production in vitro, 1534
Reovirus, gene responsible for neutralization antigen, 1305
Respiration and glycolysis, synergism between inhibitors, energy metabolism and T-cell-mediated cytolysis, 698
Reticulum cell sarcoma, nature of surface antigen inducing proliferation of normal T cells, 132
Ribonucleic acid, deoxy-, synthesis not required for activation of cytotoxic T cells by polyribonucleotides, 844

S

Sarcoma virus, avian, recovery from tumors induced by transformation-defective mutants, 1735
–, BALB/c Meth A, and unique antigen, 720
–, reticulum cell, nature of surface antigen inducing proliferation of normal T cells, 132
Sensitivity, contact, suppressor cells in tolerance, active against hapten-syngeneic and allogeneic determinants, 49
– response, contact, to picryl chloride, *I–J* subregion codes for determinants on suppressor factors, 293
Serological distinction of heavy chain variable regions of mouse immunoglobulins, common V_H determinants on heavy chains of mouse myeloma proteins with different binding sites, 1041
Serum protein β1H, requirement for cleavage of human complements C3b and C4b in solution, 257
– protein, mouse, with specific binding affinity for C4, 1001
– protein SAA-like material, amyloid-related, localization to intermediate filaments of cultured human embryonal fibroblasts, 1158
Shigella diarrhea, evidence for cell membrane toxin receptor involving β1 → 4-linked *N*-acetyl-D-glucosamine oligomers, 535
Skin and bone marrow grafts, allogeneic, long-term survival, induction of specific tissue transplantation tolerance using fractionated total lymphoid irradiation, 34
– grafts, male, pregnancy-induced unresponsiveness, adoptive transfer with thymus-dependent cells, 899
Spleen cells, normal, secondary allogeneic mixed tumor lymphocyte culture, in vitro generation of tumor-specific cytotoxic lymphocytes, 468
– cells, (responder × nonresponder) stimulated by parental macrophages, secondary antibody response in vitro to L-glutamic acid^{60}-L-alanine^{30}-L-tyrosine^{10}, 1827
Spleens, fetal, individual, specific antigen-binding cell populations, repertoire composition, size, genetic control, 394
Splenectomy, and first set rejection reactions of C57BL/6 females to male skin isografts, 809
Streptococcal antigens, group A, cross-reactive with myocardium, purification of heart-reactive antibody, isolation of streptococcal antigen, 579
– cells or cell walls, systemic injection, and arthritis, 1585
– group A-variant polysaccharide, antibody response in subjects lacking K-polypeptide

chains, 1195
Streptococcal-induced cell-mediated-immune destruction of cardiac myofibers in vitro, 344
– M-protein molecule, multiple antigenic determinants, antibody response, 1108
Suppressor cells in tolerance to contact sensitivity active against hapten-syngeneic and allogeneic determinants, 49
– factors limiting contact sensitivity response to picryl chloride, *I–J* subregion codes for determinants, 293
– T-cell activity, hapten-reactive, detection and characterization under immunization with hapten-isologous protein conjugate, 74
– T cells, hapten-reactive, inactivation by hapten-nonimmunogenic copolymers of D-amino acids, suppressor T-cell effect on helper T-cell development, 91
– T cells, neonatal, Ly phenotype and mechanism of action, 59
– T cells, specific, enrichment and surface markers, 1450
Syngeneic and allogeneic determinants, hapten-, contact sensitivity active against, suppressor cells in tolerance, 49
– cells, trinitrobenzene sulfonate-modified, cytotoxic effector cells and failure to recognize cell surface-bound trinitrophenyl dextran, 1068
– cells, trinitrophenyl- and dinitrophenyl-modified, distinguished by cytotoxic T cells, 600
– target cells, trinitrophenyl-conjugated, Ia molecules, reaction of cytotoxic T lymphocytes induced against allogeneic *I*-region determinants, 623
– viral lymphomas, *H-2*^{b} or *H-2*^{d}, and killer T lymphocytes, involvement of *H-2*^{b} or *H-2K*^{d} product in interaction, 909

T

Taenia taeniformis, vaccination using purified antigen, immunity in taeniasis-cysticercosis, 118
Taeniasis-cysticercosis immunity, vaccination against *Taenia taeniformis* using purified antigen, 118
T antigen-binding cells, inhibition by idiotypic antisera, 766
Target cells, substitution with trinitrophenol not altering induction of virus-specific modifications recognized by cytotoxic T cells, 617
– cells, syngeneic, trinitrophenyl-conjugated, Ia molecules, reaction of cytotoxic T lymphocytes induced against allogeneic *I*-region determinants, 623
T-cell activity, suppressor, hapten-reactive, detection and characterization under immunization with hapten-isologous protein conjugate, 74
–, cytotoxic, activation by polyribonucleotides, DNA synthesis not required, 844
–, cytotoxic, hemagglutinin-specific response in influenza, 893
T-cell, cytotoxic, response to ectromelia virus-infected cells, *H-2* requirement for precursor T-cell induction or lysis by effector T cells defined by BALB/c-*H-2*^{db} mutation, 869
–, helper, development, effect of suppressor T cell, and selective inactivation of hapten-reactive suppressor T cells by hapten-nonimmunogenic copolymers of D-amino acids, 91
–, helper, function, and *H-2*-linked genes, B cells expressing *Ir* genes in vitro, 1748
T-cell-mediated cytolysis, and energy metabolism, synergism between inhibitors of respiration and glycolysis, 698
T-cell mitogenesis, in man, suppression by prostaglandin, and prostaglandin-producing suppressor cell, 1719
– precursors, cytotoxic, alloantigen-specific, early postnatal development, 887
– regulation of B-cell unresponsiveness, in human peripheral blood, 1765
– subpopulations in man, help and suppression of B-cell responses by T cells bearing receptors for IgM or IgG, functional analysis, 184
–, suppressor, activity, in man, 1455
–, suppressor, activity, late-acting, antigen mediation, 1627
–, suppressor, bearing *I–J* determinants, functional and serological characterizations, 1234
–, suppressor, in mixed lymphocyte reaction in man, 368
T cells, antigen-specific, rosette-forming, nylon adherent, 1390
– bearing receptors for IgM or IgG, help and suppression of B-cell responses in man, functional analysis, 184
–, cytotoxic, cross-reactive, and expression of influenza A matrix protein on cell surface, 673
–, cytotoxic, distinguishing between trinitrophenyl- and dinitrophenyl-modified syngeneic cells, 600
–, cytotoxic, generation, and autologous mixed lymphocyte culture reactions, 1809
–, cytotoxic, recognition of virus-specific modifications, induction not altered by substitution of target cells with trinitrophenol, 617
–, effector, induction or lysis of precursor T cells, in cytotoxic T-cell response to ectromelia virus-infected cells, *H-2* requirement defined by BALB/c-*H-2*^{db} mutation, 869
–, memory, differentiation to virus plaque-forming cells and cytotoxic T lymphocytes, 11
–, normal, nature of reticulum cell sarcoma surface antigen inducing proliferation, 132
–, precursor from bone marrow, differentiation and blastogenesis, phytohemagglutinin-induced, 735
–, suppressor, generated by extract from L-glutamic acid^{50}-L-tyrosine^{50}-primed lymphoid

cells, 970
T cells, suppressor, hapten-reactive, inactivation by hapten-nonimmunogenic copolymers of D-amino acids, suppressor T-cell effect on helper T-cell development, 91
–, suppressor, neonatal, Ly phenotype and mechanism of action, 59
–, suppressor, specific, enrichment and surface markers, 1450
Thy-1.1, cell surface component, purification of milligram quantities, 986
Thymic-independent antigens, response, role of surface IgD, 1707
Thymus cell subpopulations, immunocompetent, generating cytotoxic effector cells, 412
Thymus-dependent cells, and adoptive transfer of pregnancy-induced unresponsiveness to male skin grafts, 899
Thymus-independent antigens, B-cell-activating properties, B-cell response and induction of immunological tolerance, 308
– antigens, immunological unresponsiveness, B-cell unresponsiveness to dextran, genetic mechanisms, 1663
Tissue and blood leukocytes supporting in vitro infection by dengue virus in mononuclear phagocytes, 218
– transplantation, induction of specific tolerance using fractionated total lymphoid irradiation, long-term survival of allogeneic bone marrow and skin grafts, 34
T lymphoblasts, purified mixed leukocyte culture-activated, use as autoimmunogen in inducing specific immune unresponsiveness, 1124
T-lymphocyte, murine, proliferative response, antigen presentation requiring genetic identity at major histocompatibility complex, 828
T lymphocytes bearing Ia antigens controlled by *I-J* subregion, 332
–, cytotoxic, induced against allogeneic *I*-region determinants reacting with Ia molecules on trinitrophenyl-conjugated syngeneic target cells, 623
–, cytotoxic, major histocompatibility complex (AG-B) controlling specificity, alloantigen system showing no AG-B restriction, 1773
–, cytotoxic, and virus plaque-forming cells, differentiation of memory T cells, 11
–, human, adult, role in suppressing Epstein-Barr virus in vitro, 495
–, killer, and syngeneic $H\text{-}2^b$ or $H\text{-}2^d$ viral lymphomas, involvement of $H\text{-}2^b$ or $H\text{-}2K^d$ product in interaction, 909
–, sensitized, hapten-binding receptors, lacking serological determinants of Ig constant domains, but carrying *a* locus allotypic markers, 792
–, separation of helpers and suppressors, Ly phenotypes and lack of DNA synthesis requirement for generation of concanavalin A helper and suppressor cells, 747
Tobacco glycoprotein activating factor XII, 457
Toxin receptor, cell membrane, involving $\beta 1 \rightarrow 4$-linked *N*-acetyl-D-glucosamine oligomers, in *Shigella* diarrhea, 535
Trace metals, not forming heme complexes, in regulation of heme metabolism in kidney, enzymes, 1286
Transplantation, tissue, induction of specific tolerance using fractionated total lymphoid irradiation, long-term survival of allogeneic bone marrow and skin grafts, 34
Trinitrobenzene sulfonate-modified syngeneic cells, cytotoxic effector cells and failure to recognize cell surface-bound trinitrophenyl dextran, 1068
Trinitrophenol, substitution for target cells not altering induction of virus-specific modifications recognized by cytotoxic T cells, 617
Trinitrophenyl conjugates of poly-L-(Tyr, Glu)-poly-D,L-Ala – poly-L-Lys and poly-L-(His, Glu)-poly-D,L-Ala – poly-L-Lys, primary IgM responses, *Ir* gene control, 1096
Trinitrophenyl dextran, cell surface-bound, not recognized by cytotoxic effector cells directed against trinitrobenzene sulfonate-modified syngeneic cells, 1068
Trinitrophenyl-conjugated syngeneic target cells, Ia molecules, reaction of cytotoxic T lymphocytes induced against allogeneic *I*-region determinants, 623
Trinitrophenyl- and dinitrophenyl-modified syngeneic cells distinguished by cytotoxic T cells, 600
Trinitrophenyl-modified allogeneic targets to killer cells generated by lymphocytes negatively selected to alloantigens, 361
Trypanosoma cruzi infection, immunological induction of macrophage plasminogen activator requiring thymus-derived lymphocytes, 172
– infection, modification of macrophage function, 157
Tumor, AKR, associated antigen, cell-mediated responsiveness, genetic control, locus mapped to *I* region of *H-2* complex, 1367
– cell line, monocyte, proliferation blocked by immunostimulators, 611
– virus, mammary, expression, and mammary tumors in hybrid mouse strains C57BL and GR, 1206
Tumorigenesis, radiation leukemia virus-induced, genetic control, influence of *Srlv-1*, locus not linked to *H-2*, 1088
–, radiation leukemia virus-induced, genetic control, role of major histocompatibility complex, *H-2*, 1079
Tumors induced by transformation-defective mutants, and recovery of avian sarcoma virus, 1735
–, macrophage activation, stable noncytolytic stage, 1511
–, mammary, and mammary tumor virus expression in hybrid mouse strains C57BL

and GR, 1206
poly-L-(Tyr, Glu)-poly-D,L-Ala–poly-L-Lys, trinitrophenyl conjugate, *Ir* gene control of primary IgM response, 1096
poly-L-(Tyrosine, glutamine)-poly-D,L-alanine–poly-L-lysine, antibodies produced by high and low responders, allotypic-specific analysis, 1815
L-Tyrosine^{10}, L-glutamic acid^{60}-L-alanine^{30}-secondary antibody responses in vitro by (responder × nonresponder) spleen cells stimulated by parental macrophages, 1827
L-Tyrosine^{50}, L-glutamic acid^{50}-, presence of *I-J* determinants on specific immunosuppressive factor, 287
L-Tyrosine^{50}, L-glutamic acid^{50}-, specific immunosuppressive factors, generation of suppressor T cells by extract from primed lymphoid cells, 970

V

Vaccination against *Taenia taeniformis* using purified antigen, immunity in taeniasis-cysticercosis, 118
– with vaccinia virus and immune response in man, generation of virus-specific cytotoxic activity by peripheral lymphocytes, 949
Vaccinia virus vaccination, and immune response in man, virus-specific cytotoxic activity generated by human peripheral lymphocytes, 949
Viral lymphomas, *H-2^b* or *H-2^d*, syngeneic, and killer T lymphocytes, involvement of *H-2^b* or *H-2K^d* product in interaction, 909
Virion envelope glycoprotein, and virus-induced cell-surface antigens, separate antibodies induced by immunization with syngeneic Moloney lymphoma cells, 1521
Virus antigens and histocompatibility determinants in man, effect of measles virus infection on HLA expression, 146
–, avian sarcoma, recovery from tumors induced by transformation-defective mutants, 1735
–, ectromelia, infected cells, and cytotoxic T-cell response, *H-2* requirement for precursor T-cell induction or lysis by effector T cells defined by BALB/c-*H-2^{db}* mutation, 869
–, Epstein-Barr, in vitro, role of adult human T lymphocytes in suppression, 495
–, Friend, antigenic modulation of erythroleukemic cells in vitro by serum from mice with dormant erythroleukemia, 520
–, hepatitis, blocking of in vitro and in vivo susceptibility, 1467
–, *Herpes simplex*, mitogenic properties, in interaction with lymphocytes, 1500
Virus, influenza, matrix protein antibody, detecting common antigen on surface of cells infected with influenza A viruses, 690
–, influenza A, matrix protein expression on cell surface, and recognition by cytotoxic T cells, 673
–, leukemia, AKR murine, gp71-like and BALB(X) gp71-like antigens in normal tissues in absence of overt virus expression, 422
–, leukemia, radiation, induced tumorigenesis, genetic control, influence of *Srlv-1*, locus not linked to *H-2*, 1038
–, leukemia, radiation, induced tumorigenesis, genetic control, role of major histocompatibility complex, *H-2*, 1079
–, leukemia, related antigens and alloantigens on cultured fibroblasts and sarcoma cells, unique antigen on BALB/c Meth A sarcoma, 720
–, mammary tumor, expression, and mammary tumors, in hybrid mouse strains C57BL and GR, 1206
– measles, infection in man, effect on HLA expression, 146
– plaque-forming cells and cytotoxic T lymphocytes, differentiation of memory T cells, 11
–, reo-, gene responsible for neutralization antigen, 1305
–, vaccinia, vaccination and immune response in man, virus-specific cytotoxic activity generated by human peripheral lymphocytes, 949
Viruses, dengue, and mononuclear phagocytes, enhancement by non-neutralizing antibody, 201
–, dengue, and mononuclear phagocytes, identity of blood and tissue leukocytes supporting in vitro infection, 218
–, influenza A, surface of infected cells, common antigen detected by antibody to influenza virus matrix protein, 690
Virus-induced cell surface antigens, and virion envelope glycoprotein, separate antibodies induced by immunization with syngeneic Moloney lymphoma cells, 1521
Virus-specific modifications recognized by cytotoxic T cells, induction not altered by substitution of target cells with trinitrophenol, 617

X

X-irradiation and preleukemic expression of TL antigens, 271